PATENT COOPERATION TREATY

	From the INTERNATIONAL BUREAU				
PCT	To:				
NOTIFICATION OF ELECTION	United States Patent and Trademark				
/PCT Pulo 61 2)	Office				
(PCT Rule 61.2)	(Box PCT) Crystal Plaza 2				
	Washington, DC 20231				
·	ETATS-UNIS D'AMERIQUE				
Date of mailing (day/month/year)	in its capacity as elected Office				
20 February 1998 (20.02.98)	<u> </u>				
International application No.	Applicant's or agent's file reference				
PCT/US97/10784	09010/015WO1				
International filing date (day/month/year)	Priority date (day/month/year)				
19 June 1997 (19.06.97)	19 June 1996 (19.06.96)				
Applicant					
MATHUR, Eric, J. et al					
The designated Office is hereby notified of its election man	ie.				
X in the demand filed with the International Preliminal	y Examining Authority on:				
15 January 19	98 (15.01.98)				
in a notice effecting later election filed with the International Bureau on:					
2. The election X was					
<u> </u>					
was not					
made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under					
Rule 32.2(b).					
	;				
The International Bureau of WIPO	Authorized officer				
34, chemin des Colombettes	Céline Faust				
1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Telephone No.: (41, 23) 238,83,38				
· 405/11/10 (40 (41-64) 140.14.30	Telephone No.: (41-22) 338.83.38				

PAIENT COOPERATION TREATY

PCT

NOTIFICATION CONCERNING AMENDMENTS OF THE CLAIMS

(PCT Rule 62 and Administrative Instructions, Section 417)

From the INTERNATIONAL BUREAU

To

United States Patent and Trademark Office (Box PCT) Crystal Plaza 2 Washington, DC 20231 ETATS-UNIS D'AMERIQUE

in its capacity as International Preliminary Examining Authority

Date of mailing:

20 February 1998 (20.02.98)

International application No.:

PCT/US97/10784

International filing date:

19 June 1997 (19.06.97)

Applicant:

RECOMBINANT BIOCATALYSIS, INC. et al

The International Bureau hereby informs the International Preliminary Examining Authority that no amendments under Article 19 have been received by the International Bureau (Administrative Instructions, Section 417)

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Authorised officer:

Céline Faust

Facsimile No.: (41-22) 740.14.35

Telephone No.: (41-22) 338.83.38

From the INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

To: LISA A. HAILE FISH & RICHARDSON P.C. 4225 EXECUTIVE SQUARE, SUITE 1400 LA JOLLA, CA 92037

PCT

NOTIFICATION OF TRANSMITTAL OF INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Rule 71.1)

Date of Mailing (day/month/year)

1998 APR

IMPORTANT NOTIFICATION

Applicant's or agent's file reference

09010/015W01

International filing date (day/month/year)

Priority Date (day/month/year)

PCT/US97/10784

International application No.

19 JUNE 1997

26 DECEMBER 1996

Applicant

RECOMBINANT BIOCATALYSIS, INC.

- The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international application.
- A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
- Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices. 3.

REMINDER 4.

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices)(Article 39(1))(see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide. (in Docketing Required .

RECEIVED

APR 2 3 1998

FISH & RICHARDSON, P.C.

LA JOLLA, CA

Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks

Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorize

Telephone No.

(703) 308-0196

Reviewed Hy Practice Systems

Reviewed By Billing Secretary

Immals: 🔏 8

USin for

Form PCT/IPEA/416 (July 1992)*



PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference	FOR FURTHER ACTION	See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)				
International application No.	International filing date (day/mo	nte (day/month/year) Priority date (day/month/year)				
PCT/US97/10784	19 JUNE 1997	26 DECEMBER 1996				
International Patent Classification (IPC) or national classification and IPC Please See Supplemental Sheet.						
Applicant RECOMBINANT BIOCATALYSIS, IN						
Examining Authority and is 2. This REPORT consists of a This report is also accombeen amended and are the	transmitted to the applicant act total of sheets. spanied by ANNEXES, i.e., sheet the basis for this report and/or sheet tion 607 of the Administrative In	s of the description, claims and/or drawings which have ets containing rectifications made before this Authority.				
3. This report contains indication	ns relating to the following ite	ms:				
I X Basis of the repo	nt					
II Priority						
III Non-establishme	nt of report with regard to nov	elty, inventive step or industrial applicability				
IV Lack of unity of	invention	, ,				
V X Reasoned statemen		d to novelty, inventive step or industrial applicability;				
VI Certain documents	cited					
VII Certain defects in the international application						
VIII Certain observations on the international application						
Date of submission of the demand	Date o	f completion of this report				
15 JANUARY 1998 25 FEBRUARY 1998						
Name and mailing address of the IPEAA	1	zed officer				
Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 EINAR STOVE						
Facsimile No. (703) 305-3230 Telephysia No. (703) 305-3230						
Form PCT/IPEA/409 (cover sheet) (Januar	ary 1994)*					

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.
PCT/US97/10784

	D 1 64		
I.	Basis of the report	hasis of Substitute sheets wi	hich have been furnished to the receiving Office in response to an invitation
1. T	his report has been drawn on the under Article 14 are referred to in	this report as "originally filed	d" and are not annexed to the report since they do not contain amendments):
		al application as origina	
 	X the description,	pages 1-100	, as originally filed.
	٠ ت	pages NONE	_ , filed with the demand.
		nages NONE	, filed with the letter of
		pages	, filed with the letter of
	X the claims,	Nos. 1-12	_ , as originally filed.
	x the claims,	Nos. NONE	, as amended under Article 19.
		Nos. NONE	, filed with the demand.
		Nos NONE	, filed with the letter of
		Nos	_ , filed with the letter of
	AL 20000000		, as originally filed.
	X the drawings,		, filed with the demand.
		sheets/fig NONE	, filed with the letter of
1		sheets/fig	, filed with the letter of
	x the claims, x the drawings	n established as if (some esclosure as filed, as indica	of) the amendments had not been made, since they have been considered ted in the Supplemental Box Additional observations below (Rule 70.2(c)).

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US97/10784

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement					
1. STATEMENT					
Novelty (N)	Claims		YES		
	Claims	NONE	NO		
Inventive Step (IS)	Claims Claims	I-12 NONE	YES		
Industrial Applicability (IA)	Claims		YES		
	Claims	NONE	NO		
phosphatases; 2) polynucleotides encoding or 4) thermostable alkaline phosphatases the instant claims are free of the prior and the instant claims are free	s described by SEQ rt.				

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US97/10784

Cunn	lement	1	Rav
Sunn	iement	ЖI	DUX

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of: Boxes I - VIII

Sheet 10

CLASSIFICATION:

The International Patent Classification (IPC) and/or the National classification are as listed below: IPC(6): A61K 38/46; C07H 19/00, 21/02, 21/04; C12N 9/14, 1/20, 15/00 and US Cl.: 424/94.6; 435/95, 252.3, 320.1; 536/22.1, 23.1, 23.2; 935/22

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:	} :	(11) International Publication Number: WO 97/48416
A61K 38/46, C07H 19/00, 21/02, 21/04, C12N 9/14, 1/20, 15/00	A1	(43) International Publication Date: 24 December 1997 (24.12.97)
(21) International Application Number: PCT/US (22) International Filing Date: 19 June 1997 (CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,
(30) Priority Data: 60/033,752 19 June 1996 (19.06.96) (60) Parent Application or Grant (63) Related by Continuation US 60/033,7 Filed on 19 June 1996 ((-	Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
(71) Applicant (for all designated States except US): RE NANT BIOCATALYSIS, INC. [US/US]; 505 Coa vard South, La Jolla, CA 92037 (US).	ECOMB ist Boul	
(72) Inventors; and (75) Inventors/Applicants (for US only): MATHUR, [US/US]; 2654 Galicia Way, Carlsbad, CA 920 LEE, Edd [US/US]; 2041 Paseo Dorado #4, La J 92037 (US). BYLINA, Edward [US/US]; 559 Wa Apartment A-1, Andalusia, PA 19020 (US).	009 (US Jolla, C). A
(74) Agents: HAILE, Lisa, A. et al.; Fish & Richardson P. 1400, 4225 Executive Square, La Jolla, CA 92037	.C., Sui (US).	e

(54) Title: THERMOSTABLE PHOSPHATASES

(57) Abstract

Thermostable alkaline phosphatase enzymes derived from bacteria from the genus Ammonifex, Aquifex, Archaeoglobus, Desulfuro-coccus, Methanococcus, Thermotogales, Pyrolobus, Pyrococcus, and Thermococcus organisms are disclosed. The enzymes are produced from native or recombinant host cells and can be utilized in the pharmaceutical, food, detergent, and baking industry.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	St	Slovenia
AM	Armenia	Fi	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	1E	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	1S	Iceland	MW	Malawi	US	United States of America
CA	Canada	ΙT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JР	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

THERMOSTABLE PHOSPHATASES

This invention relates to newly identified polynucleotides, polypeptides encoded by such polynucleotides, the use of such polynucleotides and polypeptides, as well as the production and isolation of such polynucleotides and polypeptides. More particularly, the polynucleotides and polypeptides of the present invention have been identified as thermostable alkaline phosphatases.

BACKGROUND OF THE INVENTION

Phosphatases are a group of enzymes that remove phosphate groups from organophosphate ester compounds. There are numerous phosphatases, including alkaline phosphatases, phosphodiesterases and phytases.

Alkaline phosphatases are widely distributed enzymes and are composed of a group of enzymes which hydrolyze organic phosphate ester bonds at alkaline pH.

Phosphodiesterases are capable of hydrolyzing nucleic acids by hydrolyzing the phosphodiester bridges of DNA and RNA. The classification of phosphodiesterases depends upon which side of the phosphodiester bridge is attacked. The 3' enzymes specifically hydrolyze the ester linkage between the 3' carbon and the phosphoric group whereas the 5' enzymes hydrolyze the ester linkage between the phosphoric group and the 5' carbon of the phosphodiester bridge. The best known of the class 3' enzymes is a phosphodiesterase from the venom of the rattlesnake or from a rustle's viper, which hydrolyses all the 3' bonds in either RNA or DNA liberating nearly all the nucleotide units as nucleotide 5' phosphates. This enzyme requires a free 3' hydroxyl group on the terminal nucleotide residue and proceeds stepwise from that end of the

polynucleotide chain. This enzyme and all other nucleases which attack only at the ends of the polynucleotide chains are called exonucleases. The 5' enzymes are represented by a phosphodiesterase from bovine spleen, also an exonuclease, which hydrolyses all the 5' linkages of both DNA and RNA and thus liberates only nucleoside 3' phosphates. It begins its attack at the end of the chain having a free 3' hydroxyl group.

Phytases are enzymes which recently have been introduced to commerce. The phytase enzyme removes phosphate from phytic acid (inositol hexaphosphoric acid), a compound found in plants such as corn, wheat and rice. The enzyme has commercial use for the treatment of animal feed, making the inositol of the phytic acid available for animal nutrition. Aspergillus ficuum and wheat are sources of phytase. (Business Communications Co., Inc., 25 Van Zant Street, Norwalk, CT 06855).

Phytase is used to improve the utilization of natural phosphorus in animal feed. Use of phytase as a feed additive enables the animal to metabolize a larger degree of its cereal feed's natural mineral content thereby reducing or altogether eliminating the need for synthetic phosphorus More important than the reduced need for additives. phosphorus additives is the corresponding reduction of phosphorus in pig and chicken waste. Many European countries severely limit the amount of manure that can be spread per acre due to concerns regarding phosphorus contamination of ground water. This is highly important in northern Europe, and will eventually be regulated throughout the remainder of the European Continent and the United States as well. (Excerpts from Business Trend Analysts, Inc., January 1994, Frost and Sullivan Report 1995 and USDA on-line information.)

Alkaline phosphatase hydrolyzes monophosphate esters, releasing an organic phosphate and the cognate alcohol compound. It is non-specific with respect to the alcohol moiety and it is this feature which accounts for the many uses of this enzyme. The enzyme has a pH optimum between 9 and 10, however, it can also function at neutral pH, (study of the enzyme industry conducted by Business Communications Company, Inc., 25 Van Zant Street, Norwalk, Connecticut 06855, 1995.).

Thermostable alkaline phosphatases are not irreversibly inactivated even when heated to 60°C or more for brief periods of time, as, for example, in the practice of hydrolyzing monophosphate esters.

Alkaline phosphatases may be obtained from numerous thermophilic organisms, such as Ammonifex degensii, Aquifex pyrophilus, Archaeoglobus lithotrophicus, Methanococcus igneus, Pyrolobus(a Crenarchaeota), Pyrococcus and Thermococcus, which are mostly Eubacteria and Euryarchaeota. Many of these organisms grow at temperatures up to about 103°C and are unable to grow below 70°C. These anaerobes are isolated from extreme environments. For example, Thermococcus CL-2 was isolated from a worm residing on a "black smoker" sulfite structure.

Interest in alkaline phosphatases from thermophilic microbes has increased recently due to their value for commercial applications. Two sources of alkaline phosphatases dominate and compete commercially: (i) animal, from bovine and calf intestinal mucosa, and (ii) bacterial, from E. coli. Due to the high turnover number of calf intestinal phosphatase, it is often selected as the label in many enzyme immunoassays. The usefulness of calf alkaline phosphatase, however, is limited by its inherently low

thermostability, which is even further compromised during the chemical preparation of the enzyme: antibody conjugates. Bacterial alkaline phosphatase is an alternative to calf alkaline phosphatase due to bacterial alkaline phosphatase's extreme thermotolerance at temperatures as high as 95°C (Tomazic-Allen, S.J., Recombinant Bacterial Phosphatase as an Immunodiagnostic Enzyme, Annals D Biology Clinique, 49(5):287-90 (1991), however, the enzyme has a very low turnover number.

There is a need for novel phosphatase enzymes having enhanced thermostability. This includes a need for enhanced alkaline phosphatases whose thermostable thermostability is beneficial in enzyme labeling processes and certain recombinant DNA techniques, such as in the dephosphorylation of vector DNA prior to insert DNA ligation. Recombinant phosphatase enzymes provide the proteins in a format amenable to efficient production of pure enzyme, which can be utilized in a variety of applications as described Accordingly, is a need there herein. characterization, amino acid sequencing, DNA sequencing, and heterologous expression of thermostable phosphatase enzymes. The present invention meets these need by providing DNA and acid sequence exprssion information and purification protocol for thermostable phosphatase derived from several organisms.

SUMMARY OF THE INVENTION

The present invention provides thermostable phosphatases from several organisms. In accordance with one aspect of the present invention, there are provided novel enzymes, as well as active fragments, analogs and derivatives thereof.

In accordance with another aspect of the present invention, there are provided isolated nucleic acid molecules

encoding the enzymes of the present invention, including mRNAs, cDNAs, genomic DNAs, as well as active analogs and fragments of such nucleic acids.

In accordance with another aspect of the present invention, there are provided isolated nucleic acid molecules encoding mature enzymes expressed by the DNA contained in the plasmid DNA vector deposited with the ATCC as Deposit No. 97536 on May 10, 1996.

In accordance with a further aspect of the present invention, there is provided a process for producing such polypeptides by recombinant techniques comprising culturing recombinant prokaryotic and/or eukaryotic host cells, containing a nucleic acid sequence of the present invention, under conditions promoting expression of said enzymes and subsequent recovery of said enzymes.

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing such enzymes for hydrolyzing monophosphate ester bonds, as an enzyme label in immunoassays, for removing 5' phosphate prior to end-labeling, and for dephosphorylating vectors prior to insert ligation.

In accordance with yet a further aspect of the present invention, there are also provided nucleic acid probes comprising nucleic acid molecules of sufficient length to specifically hybridize to a nucleic acid sequence of the present invention.

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing such enzymes, or polynucleotides encoding such enzymes, for in vitro purposes related to scientific research, for example,

to generate probes for identifying similar sequences which might encode similar enzymes from other organisms by using certain regions, i.e., conserved sequence regions of the nucleotide sequence.

These and other aspects of the present invention will be apparent to those of skill in the art from the teachings herein.

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings are illustrative of embodiments of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

Figure 1 is an illustration of the full-length-DNA and corresponding deduced amino acid sequence of Ammonifex degensii KC4 of the present invention. Sequencing was performed using a 378 automated DNA sequence for all sequences of the present invention (Applied Biosystems, Inc., Foster City, California).

Figure 2 is an illustration of the full-length DNA and corresponding deduced amino acid sequence of Methanococcus igneus Ko15.

Figure 3 is an illustration of the full-length DNA and corresponding deduced amino acid sequence of *Thermococcus alcalīphilus* AEDII12RA.

Figure 4 is an illustration of the full-length DNA and corresponding deduced amino acid sequence of *Thermococcus* celer.

PCT/US97/10784

Figure 5 is an illustration of the full-length DNA and corresponding deduced amino acid sequence of *Thermococcus* GU5L5.

Figure 6 is an illustration of the full-length DNA and corresponding deduced amino acid sequence of OC9a.

Figure 7 is an illustration of the full-length DNA and corresponding deduced amino acid sequence of M11TL.

Figure 8 is an illustration of the full-length DNA and corresponding deduced amino acid sequence of Thermococcus CL-2.

Figure 9 is an illustration of the full-length DNA and corresponding deduced amino acid sequence of $Aquife\bar{x}$ VF-5.

DETAILED DESCRIPTION OF THE INVENTION

To facilitate understanding of the invention, a number of terms are defined below.

The term "isolated" means altered "by the hand of man" from its natural state; i.e., if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a naturally occurring polynucleotide or a polypeptide naturally present in a living animal in its natural state is not "isolated", but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. For example, with respect to polynucleotides, the term isolated means that it is separated from the nucleic acid and cell in which it naturally occurs.

As part of or following isolation, such polynucleotides can be joined to other polynucleotides, such as DNAs, for mutagenesis, to form fusion proteins, and for propagation or expression in a host, for instance. The isolated polynucleotides, alone or joined to other polynucleotides such as vectors, can be introduced into host cells, in culture or in whole organisms. Introduced into host cells in culture or in whole organisms, such polynucleotides still would be isolated, as the term is used herein, because they their naturally occurring in would not be environment. Similarly, the polynucleotides and polypeptides may occur in a composition, such as a media formulation of polynucleotides introduction (solutions for polypeptides, for example, into cells or compositions or solutions for chemical or enzymatic reactions which are not naturally occurring compositions) and, therein remain isolated polynucleotides or polypeptides within the meaning of that term as it is employed herein.

The term "ligation" refers to the process of forming phosphodiester bonds between two or more polynucleotides, which most often are double stranded DNAs. Techniques for ligation are well known to the art and protocols for ligation are described in standard laboratory manuals and references, such as, for instance, Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989).

The term "oligonucleotide" as used herein is defined as a molecule comprised of two or more deoxyribonucleotides or ribonucleotides, preferably more than three, and usually more than ten. The exact size of an oligonucleotide will depend on many factors, including the ultimate function or use of the oligonucleotide. Oligonucleotides can be prepared by any suitable method, including, for example, cloning and

restriction of appropriate sequences and direct chemical synthesis by a method such as the phosphotriester method of Narang et al., 1979, Meth. Enzymol., 68:90-99; the phosphodiester method of Brown et al., 1979, Method Enzymol., 68:109-151, the diethylphosphoramidite method of Beaucage et al., 1981, Tetrahedron Lett., 22:1859-1862; the triester method of Matteucci et al., 1981, J. Am. Chem. Soc., 103:3185-3191, or automated synthesis methods; and the solid support method of U.S. Patent No. 4,458,066.

WO 97/48416

The term "plasmids" generally is designated herein by a lower case p preceded and/or followed by capital letters and/or numbers, in accordance with standard naming conventions that are familiar to those of skill in the art.

plasmids disclosed herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids by routine application of well known, published procedures. Many plasmids and other cloning and expression vectors that can be used in accordance with the present invention are well known and readily available to those of skill in the art. Moreover, those of skill readily may construct any number of other plasmids suitable for use in the invention. The properties, construction and use of such plasmids, as well as other vectors, in the present invention will be readily apparent to those of skill from the present disclosure.

The term "polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. Thus, for instance, polynucleotides as used herein refers to, among others, single-and double-stranded DNA, DNA that is a mixture of single-and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA

that may be single-stranded or, more typically, doublestranded or a mixture of single- and double-stranded regions.

In addition, polynucleotide as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide.

As used herein, the term polynucleotide includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotides" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein.

It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term polynucleotide as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including simple and complex cells, inter alia.

The term "primer" as used herein refers to an oligonucleotide, whether natural or synthetic, which is capable of acting as a point of initiation of synthesis when placed under conditions in which primer extension is initiated or possible. Synthesis of a primer extension product which is complementary to a nucleic acid strand is

initiated in the presence of nucleoside triphosphates and a polymerase in an appropriate buffer at a suitable temperature.

The term "primer" may refer to more than one primer, particularly in the case where there is some ambiguity in the information regarding one or both ends of the target region to be synthesized. For instance, if a nucleic acid sequence is inferred from a protein sequence, a "primer" generated to synthesize nucleic acid encoding said protein sequence is actually a collection of primer oligonucleotides containing sequences representing all possible codon variations based on the degeneracy of the genetic code. One or more of the primers in this collection will be homologous with the end of the target sequence. Likewise, if a "conserved" region shows significant levels of polymorphism in a population, mixtures of primers can be prepared that will amplify adjacent sequences.

The term "restriction endonucleases" and "restriction enzymes" refers to bacterial enzymes which cut doublestranded DNA at or near a specific nucleotide sequence.

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences

(introns) between individual coding segments (exons).

A coding sequence is "operably linked" to another coding sequence when RNA polymerase will transcribe the two coding sequences into a single mRNA, which is then translated into a single polypeptide having amino acids derived from both coding sequences. The coding sequences need not be

contiguous to one another so long as the expressed sequences ultimately process to produce the desired protein.

"Recombinant" enzymes refer to enzymes produced by recombinant DNA techniques; i.e., produced from cells transformed by an exogenous DNA construct encoding the desired enzyme. "Synthetic" enzymes are those prepared by chemical synthesis.

A DNA "coding sequence of" or a "nucleotide sequence encoding" a particular enzyme, is a DNA sequence which is transcribed and translated into an enzyme when placed under the control of appropriate regulatory sequences.

The term "thermostable phosphatase" refers to an enzyme which is stable to heat and heat-resistant and catalyzes the removal of phosphate groups from organophosphate ester compounds. Reference to "thermostable phosphatases" includes alkaline phosphatases, phosphodiesterases and phytases.

The phosphatase enzymes of the present invention cannot become irreversibly denatured (inactivated) when subjected to the elevated temperatures for the time necessary to effect the hydrolysis of a phosphate group from an organophosphate ester compound. Irreversible denaturation for purposes herein refers to permanent and complete loss of enzymatic activity. The phosphatase enzymes do not become irreversibly denatured from exposure to temperatures of a range from about 60°C to about 113°C or more. The extreme thermostability of the phosphatase enzymes provides additional advantages over previously characterized thermostable enzymes. Prior to the present invention, efficient hydrolysis of phosphate groups at temperatures as high as 100°C has not been demonstrated. No thermostable phosphatase has been described for this purpose.

In accordance with an aspect of the present invention, there are provided isolated nucleic acids (polynucleotides) which encode for the mature enzymes having the deduced amino acid sequences of Figures 1-9 (SEQ ID NOS:28-36).

In accordance with another aspect of the present invention, there are provided isolated polynucleotides encoding the enzymes of the present invention. The deposited material is a mixture of genomic clones comprising DNA encoding an enzyme of the present invention. Each genomic clone comprising the respective DNA has been inserted into a pBluescript vector (Stratagene, La Jolla, CA). The deposit has been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, USA, on May 10, 1996 and assigned ATCC Deposit No. 97536.

The deposit(s) have been made under the terms of the Budapest Treaty on the International Recognition of the deposit of micro-organisms for purposes of patent procedure. The strains will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. These deposits are provided merely as convenience to those of skill in the art and are not an admission that a deposit be required under 35 U.S.C. §112. The sequences of the polynucleotides contained in the deposited materials, as well as the amino acid sequences of the polypeptides encoded thereby, are controlling in the event of any conflict with any description of sequences herein. A license may be required to make, use or sell the deposited materials, and no such license is hereby granted.

The polynucleotides of this invention were originally recovered from genomic gene libraries derived from the following organisms:

Ammonifex degensii KC4 is a eubacteria from the genus Ammonifex. It was isolated in Java, Indonesia. It is a gram-negative, chemolithoautotroph. It grows optimally at $70\,^{\circ}\text{C}$ in a low-salt culture medium at pH 7 with 0.2% nitrate as a substrate and H_2/CO_2 in gas phase.

Methanococcus igneus KOL5 is a Euryarchaeota isolated from Kolbeinsey Ridge in the north of Iceland. It grows optimally at 85°C and pH 7.0 in a high-salt marine medium with $\rm H_2/CO_2$ in a gas phase. Aquifex pyrophilus KOL 5A is a marine bacteria isolated from th Kolbeinsey Ridge in the north of Iceland. It is a gram-negative, rod-shaped, strictly chemolithoautotrophic, knall gas bacterium, and a denitrifier. It grows optimally at 85°C in high-salt marine medium at pH 6.8 with $\rm O_2$ as a substrate and $\rm H_2/CO_2 + 0.5\% O_2$ in gas phase.

Thermococcus alcaliphilus AEDII12RA is from the genus Thermococcus. AEDII12RA grows optimally at 85°C, pH 9.5 in a high salt medium (marine) containing polysulfides and yeast extract as substrates and N_2 in gas phase.

Thermococcus celer is an Euryarchaeota. It grows optimally at $85\,^{\circ}\text{C}$ and pH 6.0 in a high-salt marine medium containing elemental sulfur, yeast extract, and peptone as substrates and N_2 in gas phase.

Thermococcus GU5L5 is an Euryarchaeota isolated from the Guaymas Basin in Mexico. It grows optimally at 85°C and pH 6.0 in a high-salt marine medium containing 1% elemental sulfur, 0.4% yeast extract, and 0.5% peptone as substrates with N_2 in gas phase.

OC9a-27A3A is a bacteria of unknown etilogy obtained from Yellowstone National Park and maintained as a pure



culture. It grows well on a TK6 medium and has cellulose degrader activity. Further, it codes for an alkaline phosphatase having greater than 50% polypeptide identity and greater than 32% polynucleotide identity to each of Bombyx mori and Escherichia coli C alkaline phosphatase precursors, which is significant homologyy. Thus, it is expected that OC9a-27A3A can be cloned and expressed readily in Escherichi Coli C in place of its native alkaline phosphatase precursor.

Mil TL is a new species of Desulfurococcus isolated from Diamond Pool in Yellowstone National Park. MilTL grows heterotrophically by fermentation of different organic materials (sulfur is not necessary) and forms grape-like aggregates. The organism grows optimally at 85°C to 88°C and pH 7.0 in a low salt medium containing yeast extract, peptone, and gelatin as substrates with an N_2/CO_2 gas phase.

Thermococcus CL-2 is an Euryarchaeota isolated from the North Cleft Segment in the Juan de Fuca Ridge. It grows optimally at 88°C in a salt medium with an argon atmosphere.

Aquifex VF-5 is a marine bacteria isolated from a beach in Vulcano, Italy. It is a gram-negative, rod-shaped, strictly chemolithoautotrophic, knall gas bacterium. It grows optimally from 85-90°C in high-salt marine medium at pH 6.8, with O_2 as a substrate and H_2/CO_2 + 0.5% O_2 in gas phase.

Accordingly, the polynucleotides and enzymes encoded thereby are identified by the organism from which they were isolated, and are sometimes hereinafter referred to as "KC4" (Figure 1 and SEQ ID NOS:19 and 28), "Ko15" (Figure 2 and SEQ ID NOS:20 and 29), "AEDII12RA" (Figure 3 and SEQ ID NOS:21 and 30), "Celer" (Figure 4 and SEQ ID NOS:22 and 31), "GU5L5" (Figure 5 and SEQ ID NOS:23 and 32), "OC9a" (Figure 6 and SEQ ID NOS:24 and 33), "M11TL" (Figure 7 and SEQ ID NOS:25 and

34), "CL-2" (Figure 8 and SEQ ID NOS:26 and 35) and "VF-5" (Figure 9 and SEQ ID NOS:27 and 36).

The polynucleotides and polypeptides of the present invention show identity of the nucleotide and protein level to known genes and proteins encoded thereby as shown in Table 1.

Table 1

Clone	Gene/Protein with Closest Homology	Protein Identity	Nucleic Acid Identity
Ammonifex degensiii KC4-3A1A	Yarrowia lipolytica, Candida lipolytica, acid phosphatase	47%	24%
Ammonifex degensii KC4-3A1A	Saccharomyces cerevisiae, hypothetical protein YBR094w	54%	26%
Methanococcus igeneus Kol5-9A1A	Yarrowia lipolytica, Candida lipolytica, acid phosphatase	45%	25%
Methanococcus igeneus Kol5-9A1A	Saccharomyces cerevisiae, hypothetical protein YBR094w, hypothetical protein YBR0821	52%	25%
Thermococcus alcaliphilus AEDII12RA-18A	No homology found		
Thermococus celer 25A1A	No homology found		
Thermococcus GU5L5- 26A1A	Bacillius subtilis, alkaline phosphatase IV precursor, alkaline phosphomonoesterase, glycerophosphatase, and phosphomonoesterase	58%	38%
Thermococcus GU5L5- 26A1A	Bacillius subtilis, alkaline phosphatase III precursor	58%	37%
OC9a-27A3A	Bombyx mori (silkworm), alkaline phosphatase precursor	54%	33 %
OC9a - 27A3A	Escherichia coli C, alkaline phosphatase precursor	53 %	34%
M11 TL - 29A1A	Rhodobacter capsulatus, hypothetical protein B	43 %	24%
Thermococcus C12-30A1A	Yarrowia lipolytica, Candida lipolytica, acid phosphatase	49%	27%
Thermococcus CL2-30A1A	Saccharomyces cerevisiae, hypothetical protein YBR094w hypothetical protein YBR0821	50%	25%
Aquifex VF5-34A1A	Escherichia coli, suppressor protein suhB	57%	34%

All of the clones identified in Table 1 encode polypeptides which have phosphatase activity.

One means for isolating the nucleic acid molecules encoding the enzymes of the present invention is to probe a gene library with a natural or artificially designed probe using art recognized procedures (see, for example: Current Protocols in Molecular Biology, Ausubel F.M. et al. (EDS.) Green Publishing Company Assoc. and John Wiley Interscience, New York, 1989, 1992). It is appreciated by one skilled in the art that the polynucleotides of SEQ ID NOS: 1-18, or fragments thereof (comprising at least 12 contiguous nucleotides), are particularly useful probes. Other particularly useful probes for this purpose are hybridizable fragments of the sequences of SEQ ID NOS: 19-27 (i.e., comprising at least 12 contiguous nucleotides).

With respect to nucleic acid sequences which hybridize specific nucleic acid sequences disclosed herein, hybridization may be carried out under conditions of reduced stringency, medium stringency or even stringent conditions. As an example of oligonucleotide hybridization, a polymer membrane containing immobilized denatured nucleic acids is first prehybridized for 30 minutes at 45°C in a solution consisting of 0.9 M NaCl, 50 mM NaH₂PO₄, pH 7.0, 5.0 mM 0.5% SDS, 10X Denhardt's, and 0.5 Na₂EDTA, polyriboadenylic acid. Approximately 2 X 107 cpm (specific activity 4-9 X 108 cpm/ug) of 32P end-labeled oligonucleotide probe are then added to the solution. After 12-16 hours of incubation, the membrane is washed for 30 minutes at room temperature in 1X SET (150 mM NaCl, 20 mM Tris hydrochloride, pH 7.8, 1 mM Na₂EDTA) containing 0.5% SDS, followed by a 30 minute wash in fresh 1X SET at (Tm less 10°C) for the oligonucleotide probe. The membrane is then exposed to autoradiographic film for detection of hybridization signals.

Stringent conditions means hybridization will occur only if there is at least 90% identity, preferably at least 95% identity and most preferably at least 97% identity between the sequences. Further, it is understood that a section of a 100 bps sequence that is 95 bps in length has 95% identity with the 1090 bps sequence from which it is obtained. See J. Sambrook et al., Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory (1989) which is hereby incorporated by reference in its entirety. Also, it is understood that a fragment of a 100 bps sequence that is 95 bps in length has 95% identity with the 100 bps sequence from which it is obtained.

As used herein, a first DNA (RNA) sequence is at least 70% and preferably at least 80% identical to another DNA (RNA) sequence if there is at least 70% and preferably at least a 80% or 90% identity, respectively, between the bases of the first sequence and the bases of the another sequence, when properly aligned with each other, for example when aligned by BLASTN.

The present invention relates to polynucleotides which differ from the reference polynucleotide such that the differences are silent, for example, the amino acid sequence encoded by the polynucleotides is the same. The present invention also relates to nucleotide changes which result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference polynucleotide. In a preferred aspect of the invention these polypeptides retain the same biological action as the polypeptide encoded by the reference polynucleotide.

The polynucleotides of this invention were recovered from genomic gene libraries from the organisms listed in Table 1. Gene libraries were generated from either of a

Lambda ZAP II or a pBluscript] cloning vector (Stratagene Cloning Systems). Mass excisions were performed on these libraries to generate libraries in the pBluescript phagemid. Libraries were generated and excisions were performed according to the protocols/methods hereinafter described.

The polynucleotides of the present invention may be in the form of RNA or DNA which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The coding sequences which encodes the mature enzymes may be identical to the coding sequences shown in Figures 1-9 (SEQ ID NOS: 19-27) or may be a different coding sequence which coding sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same mature enzymes as the DNA of Figures 1-9 (SEQ ID NOS: 19-27).

The polynucleotide which encodes for the mature enzyme of Figures 1-9 (SEQ ID NOS: 28-36) may include, but is not limited to: only the coding sequence for the mature enzyme; the coding sequence for the mature enzyme and additional coding sequence such as a leader sequence or a proprotein sequence; the coding sequence for the mature enzyme (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature enzyme.

Thus, the term "polynucleotide encoding an enzyme (protein)" encompasses a polynucleotide which includes only coding sequence for the enzyme as well as a polynucleotide which includes additional coding and/or non-coding sequence.

The present invention further relates to variants of the hereinabove described polynucleotides which encode for

fragments, analogs and derivatives of the enzymes having the deduced amino acid sequences of Figures 1-9 (SEQ ID NOS: 28-36). The variant of the polynucleotide may be a naturally occurring allelic variant of the polynucleotide or a non-naturally occurring variant of the polynucleotide.

Thus, the present invention includes polynucleotides encoding the same mature enzymes as shown in Figures 1-9 (SEQ ID NOS: 19-27) as well as variants of such polynucleotides which variants encode for a fragment, derivative or analog of the enzymes of Figures 1-9 (SEQ ID NOS: 19-27). Such nucleotide variants include deletion variants, substitution variants and addition or insertion variants.

As hereinabove indicated, the polynucleotides may have a coding sequence which is a naturally occurring allelic variant of the coding sequences shown in Figures 1-9 (SEQ ID NOS: 19-27). As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded enzyme. Also, using directed and other evolution strategies, one may make very minor changes in DNA sequence which can result in major changes in function.

Fragments of the full length gene of the present invention may be used as hybridization probes for a cDNA or a genomic library to isolate the full length DNA and to isolate other DNAs which have a high sequence similarity to the gene or similar biological activity. Probes of this type preferably have at least 10, preferably at least 15, and even more preferably at least 30 bases and may contain, for example, at least 50 or more bases. In fact, probes of this type having at least up to 150 bases or greater may be preferably utilized. The probe may also be used to identify

a DNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete gene including regulatory and promotor regions, exons and introns. An example of a screen comprises isolating the coding region of the gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary or identical to that of the gene or portion of the gene sequences of the present invention are used to screen a library of genomic DNA to determine which members of the library the probe hybridizes to.

It is also appreciated that such probes can be and are preferably labeled with an analytically detectable reagent to facilitate identification of the probe. Useful reagents include but are not limited to radioactivity, fluorescent dyes or enzymes capable of catalyzing the formation of a detectable product. The probes are thus useful to isolate complementary copies of DNA from other sources or to screen such sources for related sequences.

further relates invention present polynucleotides which hybridize to the hereinabove-described sequences if there is at least 70%, preferably at least 90%, and more preferably at least 95% identity between the sequences. (As indicated above, 70% identity would include within such definition a 70 bps fragment taken from a 100 bp polynucleotide, for example.) The present invention particularly relates to polynucleotides which hybridize under hereinabove-described conditions the to stringent As herein used, the term "stringent polynucleotides. conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the The polynucleotides which hybridize to the hereinabove described polynucleotides in a preferred embodiment encode enzymes which either retain substantially

the same biological function or activity as the mature enzyme encoded by the DNA of Figures 1-9 (SEQ ID NOS: 19-27). In referring to identity in the case of hybridization, as known in the art, such identity refers to the complementarity of two polynucleotide segments.

Alternatively, the polynucleotide may have at least 15 bases, preferably at least 30 bases, and more preferably at least 50 bases which hybridize to any part of a polynucleotide of the present invention and which has an identity thereto, as hereinabove described, and which may or may not retain activity. For example, such polynucleotides may be employed as probes for the polynucleotides of SEQ ID NOS: 19-27, for example, for recovery of the polynucleotide or as a diagnostic probe or as a PCR primer.

Thus, the present invention is directed to polynucleotides having at least a 70% identity, preferably at least 90% identity and more preferably at least a 95% identity to a polynucleotide which encodes the enzymes of SEQ ID NOS: 28-36 as well as fragments thereof, which fragments have at least 15 bases, preferably at least 30 bases, more preferably at least 50 bases and most preferably fragments having up to at least 150 bases or greater, which fragments are at least 90% identical, preferably at least 95% identical and most preferably at least 97% identical to any portion of a polynucleotide of the present invention.

The present invention further relates to enzymes which have the deduced amino acid sequences of Figures 1-9 (SEQ ID NOS: 28-36) as well as fragments, analogs and derivatives of such enzyme.

The terms "fragment," "derivative" and "analog" when referring to the enzymes of Figures 1-9 (SEQ ID NOS. 28-36)

means enzymes which retain essentially the same biological function or activity as such enzymes. Thus, an analog includes a proprotein which can be activated by cleavage of the proprotein portion to produce an active mature enzyme.

The enzymes of the present invention may be a recombinant enzyme, a natural enzyme or a synthetic enzyme, preferably a recombinant enzyme.

The fragment, derivative or analog of the enzymes of Figures 1-9 (SEQ ID NOS.28-36) may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature enzyme is fused with another compound, such as a compound to increase the half-life of the enzyme (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature enzyme, such as a leader or secretory sequence or a sequence which is employed for purification of the mature enzyme or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

The enzymes and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

The term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or enzyme present in a living animal

is not isolated, but the same polynucleotide or enzyme, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or enzymes could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

The enzymes of the present invention include the enzymes of SEQ ID NOS: 28-36 (in particular the mature enzyme) as well as enzymes which have at least 70% similarity (preferably at least 70% identity) to the enzymes of SEQ ID NOS: 28-36 and more preferably at least 90% similarity (more preferably at least 90% identity) to the enzymes of SEQ ID NOS: 28-36 and still more preferably at least 95% similarity (still more preferably at least 95% identity) to the enzymes of SEQ ID NOS: 28-36 and also include portions of such enzymes with such portion of the enzyme generally containing at least 30 amino acids and more preferably at least 50 amino acids and most preferably at least up to 150 amino acids.

As known in the art "similarity" between two enzymes is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one enzyme to the sequence of a second enzyme. The definition of 70% similarity would include a 70 amino acid sequence fragment of a 100 amino acid sequence, for example, or a 70 amino acid sequence obtained by sequentially or randomly deleting 30 amino acids from the 100 amino acid sequence.

A variant, i.e. a "fragment", "analog" or "derivative" polypeptide, and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions, fusions and truncations, which may be present in any combination.

Among preferred variants are those that vary from a reference by conservative amino acid substitutions. Such substitutions are those that substitute a given amino acid in a polypeptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile; interchange of the hydroxyl residues Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg and replacements among the aromatic residues Phe, Tyr.

Most highly preferred are variants which retain the same biological function and activity as the reference polypeptide from which it varies.

Fragments or portions of the enzymes of the present invention may be employed for producing the corresponding full-length enzyme by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length enzymes. Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

The present invention also relates to vectors which include polynucleotides of the present invention, host cells which are genetically engineered with vectors of the invention and the production of enzymes of the invention by recombinant techniques.

Host cells are genetically engineered (transduced or transformed or transfected) with the vectors of this invention which may be, for example, a cloning vector such as an expression vector. The vector may be, for example, in the

form of a plasmid, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the genes of the present invention. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

The polynucleotides of the present invention may be employed for producing enzymes by recombinant techniques. Thus, for example, the polynucleotide may be included in any one of a variety of expression vectors for expressing an enzyme. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other vector may be used as long as it is replicable and viable in the host.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the $E.\ coli.\ lac$ or trp, the phage lambda P_L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses.

The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

The vector containing the appropriate DNA sequence as hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein.

As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as E. coli, Streptomyces, Bacillus subtilis; fungal cells, such as yeast; insect cells such as Drosophila S2 and Spodoptera Sf9; animal cells such as CHO, COS or Bowes melanoma; adenoviruses; plant cells, etc. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors

and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example; Bacterial: pQE70, pQE60, pQE-9 (Qiagen), pBluescript II KS, ptrc99a, pKK223-3, pDR540, pRIT2T. (Pharmacia); Eukaryotic: pXT1, pSG5 (Stratagene) pSVK3, pBpV, pMSG, pSVL SV40 (Pharmacia). However, any other plasmid or vector may be used as long as they are replicable and viable in the host.

Promoter regions can be selected from any desired gene using CAT (chloramphenical transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda $P_{\rm R}$, P and trp. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis, L., Dibner, M., Battey, I., Basic Methods in Molecular Biology, (1986)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the enzymes of the

invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), the disclosure of which is hereby incorporated by reference.

Transcription of the DNA encoding the enzymes of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of E. coli and S. cerevisiae TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences,

and preferably, a leader sequence capable of directing secretion of translated enzyme. Optionally, the heterologous sequence can encode a fusion enzyme including an N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product.

Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include Et coli, Bacillus subtilis, Salmonella typhimurium and various species within the genera Pseudomonas, Streptomyces, and Staphylococcus, although others may also be employed as a matter of choice.

As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1-(Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g.,

temperature shift or chemical induction) and cells are cultured for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well known to those skilled in the art.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell, 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

The enzyme can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used,

as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The enzymes of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the enzymes of the present invention may be glycosylated or may be non-glycosylated. Enzymes of the invention may or may not also include an initial methionine amino acid residue.

Phosphatases are a group of key enzymes in the removal of phosphate groups from organophosphate ester compounds. There are numerous phosphatases, including alkaline phosphatases, phosphodiesterases and phytases.

The general application and definitions of such compounds are discussed above under the background of the invention section.

The present invention provides novel phosphatase enzymes having enhanced thermostability. Such phosphatases are beneficial in enzyme labeling processes and in certain recombinant DNA techniques, such as in the dephosphorylation of vector DNA prior to insert DNA ligation. The recombinant phosphatase enzymes provide the proteins in a format amenable to efficient production of pure enzyme, which can be utilized in a variety of applications as described herein.

Antibodies generated against the enzymes corresponding to a sequence of the present invention can be obtained by direct injection of the enzymes into an animal or by

administering the enzymes to an animal, preferably a nonhuman. The antibody so obtained will then bind the enzymes itself. In this manner, even a sequence encoding only a fragment of the enzymes can be used to generate-antibodies binding the whole native enzymes. Such antibodies can then be used to isolate the enzyme from cells expressing that enzyme.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, Nature, 256:495-497, 1975), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today 4:72, 1983), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96, 1985).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic enzyme products of this invention. Also, transgenic mice may be used to express humanized antibodies to immunogenic enzyme products of this invention.

Antibodies generated against an enzyme of the present invention may be used in screening for similar enzymes from other organisms and samples. Such screening techniques are known in the art, for example, one such screening assay is described in Sambrook and Maniatis, Molecular Cloning: A Laboratory Manual (2d Ed.), vol. 2:Section 8.49, Cold Spring Harbor Laboratory, 1989, which is hereby incorporated by reference in its entirety.

The present invention will be further described with reference to the following examples; however, it is to be understood that the present invention is not limited to such examples. All parts or amounts, unless otherwise specified, are by weight.

In order to facilitate understanding of the following examples certain frequently occurring methods and/or terms will be described.

"Plasmids" are designated by a lower case "p" preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan.

"Digestion" of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. analytical purposes, typically 1 μ g of plasmid or fragment is used with about 2 units of enzyme in about 20 μ l of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 μg of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is

electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Size separation of the cleaved fragments is performedusing 8 percent polyacrylamide gel described by Goeddel et al., Nucleic Acids Res., 8:4057 (1980).

"Oligonucleotides" refers to either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands which may be chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide will ligate to a fragment that has not been dephosphorylated.

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments (Maniatis, T., et al., Id., p. 146). Unless otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units of T4 DNA ligase ("ligase") per 0.5 μ g of approximately equimolar amounts of the DNA fragments to be ligated.

Unless otherwise stated, transformation was performed as described in Sambrook and Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, 1989.

One means for isolating the nucleic acid molecules encoding the enzymes of the present invention is to probe a gene library with a natural or artificially designed probe using art recognized procedures (see, for example: Current Protocols in Molecular Biology, Ausubel F.M. et al. (EDS.) Green Publishing Company Assoc. and John Wiley Interscience,

New York, 1989, 1992). It is appreciated to one skilled in the art that the polynucleotides of SEQ ID NOS:1-16, or fragments thereof (comprising at least 10 or 12 contiguous nucleotides), are particularly useful probes. Other particularly useful probes for this purpose are fragments hybridizable fragments to the sequences of SEQ ID NOS:19-27 (i.e., comprising at least 10 or 12 contiguous nucleotides).

It is also appreciated that such probes can be and are preferably labeled with an analytically detectable reagent to facilitate identification of the probe. Useful reagents include but are not limited to radioactivity, fluorescent dyes or enzymes capable of catalyzing the formation of a detectable product. The probes are thus useful to isolate complementary copies of DNA from other sources or to screen such sources for related sequences.

With respect to nucleic acid sequences which hybridize specific nucleic acid sequences disclosed herein, hybridization may be carried out under conditions of reduced stringency, medium stringency or even stringent conditions. As an example of oligonucleotide hybridization, a polymer membrane containing immobilized denatured nucleic acids is first prehybridized for 30 minutes at 45°C in a solution consisting of 0.9 M NaCl, 50 mM NaH, PO4, pH 7.0, 5.0 mM 0.5% SDS, 10X Denhardt's, and 0.5 Na, EDTA, polyriboadenylic acid. Approximately 2 X 10' cpm (specific activity 4-9 X 108 cpm/ug) of 32P end-labeled oligonucleotide probe are then added to the solution. After 12-16 hours of incubation, the membrane is washed for 30 minutes at room temperature in 1X SET (150 mM NaCl, 20 mM Tris hydrochloride, pH 7.8, 1 mM Na₂EDTA) containing 0.5% SDS, followed by a 30 minute wash in fresh 1X SET at Tm -10°C for the oligo-

nucleotide probe. The membrane is then exposed to autoradiographic film for detection of hybridization signals.

Stringent conditions means hybridization will occur only if there is at least 90% identity, preferably 95% identity and most preferably at least 97% identity between the sequences. See J. Sambrook et al., Molecular Cloning, A Laboratory Manual (2d Ed. 1989) (Cold Spring Harbor Laboratory) which is hereby incorporated by reference in its entirety.

"Identity" as the term is used herein, refers to a polynucleotide sequence which comprises a percentage of the same bases as a reference polynucleotide (SEQ ID NOS:1-16). For example, a polynucleotide which is at least 90% identical to a reference polynucleotide, has polynucleotide bases which are identical in 90% of the bases which make up the reference polynucleotide and may have different bases in 10% of the bases which comprise that polynucleotide sequence.

The present invention relates to polynucleotides which differ from the reference polynucleotide such that the differences are silent changes, for example, the amino acid sequence encoded by both polynucleotides is the same. The present invention also relates to nucleotide changes which result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference polynucleotide. In a preferred aspect of the invention these polypeptides retain the same biological action as the polypeptide encoded by the reference polynucleotide.

The polynucleotides of this invention were recovered from genomic gene libraries from the organisms listed in Table 1. Gene libraries were generated in the Lambda ZAP II

cloning vector (Stratagene Cloning Systems). Mass excisions were performed on these libraries to generate libraries in the pBluescript phagemid. Libraries were generated and excisions were performed according to the protocols/methods hereinafter described.

The excision libraries were introduced into the *E. coli* strain BW14893 F'kanlA. Expression clones were then identified using a high temperature filter assay using phosphatase buffer containing 1 mg/ml BCIP (5-Bromo-4-chloro-3-indolyl phosphate). Expression clones encoding BCIPases were identified and repurified from the following organisms: Ammonifex degensii KC4, Methanococcus igneus KoL5, Thermococcus alcaliphilus AED112RA, Thermococcus celer, Thermococcus GU5L5, OC9a, M11TL, Thermococcus CL-2 and Aquifex VF-5.

Expression clones were identified by use of a high temperature filter assay with either acid phosphatase buffer or alkaline phosphatase buffer containing BCIP. Metcalf, et al., Evidence for two phosphonate degradative pathways in Enterobacter Aerogenes, J. Bacteriol., 174:2501-2510 (1992)).

BCIPase activity was tested as follows: An excision library was introduced into the $E.\ Coli$ strain BW14893 F'kan, a pho'pnh'lac' strain. After growth on 100 mm LB plates containing 100 $\mu g/ml$ ampicillin, 80 $\mu g/ml$ methicillin and 1mM IPTG, colony lifts were performed using Millipore HATF membrane filters. The colonies transferred to the filters were lysed with chloroform vapor in 150 mm glass petri dishes. The filters were transferred to 100 mm glass petri dishes containing a piece of Whatman 3MM filter paper saturated with either acid phosphatase buffer (see recipe below) or alkaline phosphatase buffer (see recipe below) containing no BCIP. The dish was placed in the oven at 80-

85°C for 30-45 minutes to heat inactivate endogenous *E. coli* phosphatases. The filter bearing lysed colonies were then transferred to a 100 mm glass petri dish containing 3MM paper saturated with either acid phosphatase buffer or alkaline phosphatase buffer containing 1 mg/ml BCIP. The dish was placed in the oven at 80-85°C.

Alkaline Phosphatase Buffer (referenced in Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, p. 1874) includes 100 mM NaCl, 5 mM MgCl₂ and 100 mM Tris-HCl (pH 9.5). Clones expressing phosphatase activity (when the alkaline phosphatase buffer was used) were derived from libraries derived from the organism identified above.

Acid Phosphatase Buffer includes 100 mM NaCl, 5 mM MgCl $_2$ and 100 mM Tris-HCL (pH 6.8). Clones expressing phosphatase activity (when the acid phosphatase buffer was used) were derived from the library derived from MllTL.

'Positives' were observed as blue spots on the filter membranes. The following filter rescue technique was used to retrieve plasmid from lysed positive colony.

Filter Rescue Technique: A pasteur pipette (or glass capillary tube) was used to core blue spots on the filter membrane. The small filter disk was placed in an Eppendorf tube containing 20 ul of deionized water. The Eppendorf tube was incubated at 75°C for 5 minutes followed by vortexing to elute plasmid DNA off the filter. Plasmid DNA containing DNA inserts from Thermococcus alcaliphilus AEDII12RA was used to DH10B electrocompetent E . coli transform Electrocompetent BW14893 F'kanlA E. coli cells were used for transformation of plasmid DNA containing inserts from Ammonifex degensii KC4, Methanococcus igneus KOL5, and Thermococcus GU5L5. The filter-lift assay was repeated on

transformation plates to identify 'positives.' The transformation plates were returned to 37°C incubator to regenerate colonies. 3 ml of LBamp liquid was inoculated with repurified positives and incubated at 37°C overnight. Plasmid DNA was isolated from these cultures and plasmid insert were sequenced.

In some instances where the plates used for the initial colony lifts contained non-confluent colonies, a specific colony corresponding to a blue spot on the filter could be identified on a regenerated plate and repurified directly, instead of using the filter rescue technique. This "repurification" protocol was used for plasmid DNA containing inserts from the following: Ammonifex degensii KC4, Thermococcus celer, M11TL, and Aquifex VF-5.

The filter rescue technique was used for DNA from the following organisms: Ammonifex degensii KC4, Methanococcus igneus KOL5, Thermococcus alcaliphilus AED1112RA, Thermococcus CL-2, and OC9a.

Phosphatases are a group of key enzymes that remove phosphate groups from organophosphate ester compounds. The most important phosphatases for commercial purposes are alkaline phosphatases, phosphodiesterases, and phytases.

Alkaline phosphatases have several commercial applications, including their use in analytical applications as an_enzyme label in ELISA immunoassays and enzyme-linked gene probes, and their use in research applications for removing 5' phosphates in polynucleotides prior to end-labeling and for dephosphorylating vectors prior to insert ligation (see also Current Protocols in Molecular Biology, (John Wiley & Sons) (1995), chapter 3, section 10).

Alkaline phosphatase hydrolyzes monophosphate esters, releasing inorganic phosphate and the cognate alcohol compound. It is non-specific with respect to the alcohol moiety, a feature which accounts for the many uses of this enzyme. The enzyme has a pH optimum between 9 and 10, however, it can also work at neutral pH. (From a study of the enzyme industry conducted by Business Communications, Co., Inc., 25 Van Zant Street, Norwalk, CT 06855, 1995.)

Two sources of alkaline phosphatase dominate and compete animal, from bovine and calf intestinal in the market: mucosa, and bacterial, from E. coli. Due to the high turnover number of calf intestinal phosphatase, it is often selected as the label in many enzyme immunoassays. usefulness of calf alkaline phosphatase is limited by its inherently low thermal stability, which is even further compromised during the chemical preparation of enzyme: antibody conjugates. Bacterial alkaline phosphatase could be an attractive alternative to calf alkaline phosphatase due to bacterial alkaline phosphatase's extreme thermotolerance at temperatures as high as 95°C. (Tomazic-Allen S.J., Recombinant bacterial alkaline phosphatase as immunodiagnostic enzyme, Annales de Biologie Clinique, 1991, 49(5):287-90).

Antibodies generated against the enzymes corresponding to a sequence of the present invention can be obtained by direct injection of the enzymes into an animal or by adminIstering the enzymes to an animal, preferably a nonhuman. The antibody so obtained will then bind the enzymes itself. In this manner, even a sequence encoding only a fragment of the enzymes can be used to generate antibodies binding the whole native enzymes. Such antibodies can then be used to isolate the enzyme from cells expressing that enzyme.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, 1975, Nature, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, Immunology Today 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole, et al., 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic enzyme products of this invention. Also, transgenic mice may be used to express humanized antibodies to immunogenic enzyme products of this invention.

Antibodies, as described above, may be employed as a probe to screen a library to identify the above-described activities or cross-reactive activities in gene libraries generated from the organisms described above or other organisms.

Example 1

Bacterial Expression and Purification of Alkaline Phosphatase Enzymes

DNA encoding the enzymes of the present invention, SEQ ID NOS:1 through 16, were initially amplified from a pBluescript vector containing the DNA by the PCR technique using the primers noted herein. The amplified sequences were then inserted into the respective pQE vector listed beneath the primer sequences, and the enzyme was expressed according to the protocols set forth herein. The 5' and 3' oligonucleotide primer sequences used for subcloning and vectors for the respective genes are as follows:

Ammonifex degensii KC4 - 3A1A

5' CCGA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG GGG GCA GGT CCG AAA AGG 3' 5' CCGA GGA TCC TCA CCG CCC CCT GCG GGT GCG 3' Vector: pQET3

Methanococcus igneus Kol5 - 9A1A

5' CCGA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG TTG GAT ATA CTG CTT GTT 3' 5' CCGA CGA TCC TTA TTT TTT AAC CAA ATGT TCC 3' PQET3

Thermococcus Alcaliphilus AEDII12RA -18A

5' CCGA CAA TTG ATT AAA GAG GAG AAA TTA ACT ATG ATG ATG GAA TTC ACT CGC 3' 5' CGGA GGA TCC CTA CAG TTC TAA AAG TCT TTT A 3' Vector: pQET3

Thermococcus Celer 25AlA (incorporating Mfel restriction site)

5' CCGA CAA TTG ATT AAA GAG GAG AAA TTA ACT ATG AGA ACC CTG ACA ATA AAC 3' 5' CCGA GGA TCC TTA CAC CCA CAG AAC CCT TAC 3' Vector pQET3

Thermococcus GU5L5 - 26A1A

5' CCGA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG AAA GGA AAG TCT CTT GTT 3' 5' CCGA GGA TCC TCA AGC TTC CTG GAG AAT CAA 3' Vector pQET3

OC9a - 27A3A

5' CCGA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG CCA AGA AAT ATG GCC GCT 3' 5' CCGA GGA TCC TTA AGG CTT CTC GAG GTG GGG GTT 3' $\tt Vector\ pQET3$

M11 TL - 29A1A (incorporating Mfel restriction site)

5' CCGA CAA TTG ATT AAA GAG GAG AAA TTA ACT ATG TAT AAA TGG ATT ATT GAG GG 3' 5' CCGA GGA CTA AAC ATA GTC TAA GTA ATT AGC 3' Vector pQET3

Thermococcus CL-2 - 30A1A

5' CCGA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG AGA ATC CTC CTC ACC AAC 3' 5' CCGA GGA TCC TCA CAG GCT CAG AAG CCT TTG 3' Vector pQET3

Aquifex VF-5 - 34A1A

5' CCGA GAA TTC ATT AAA GAG GAG AAA TTA ACT ATG GAA AAC TTA AAA AAG TAC CT 3' 5' CCGA GGA TCC TCA CCG CCC CCT GCG GGT GCG 3' Vector pQET3

The restriction enzyme sites indicated correspond to the restriction enzyme sites on the bacterial expression vector indicated for the respective gene (Qiagen, Inc. Chatsworth, CA). The pQE vector encodes antibiotic resistance (Amp $^{\rm r}$), a bacterial origin of replication (ori), an IPTG-regulatable promoter operator (P/O), a ribosome binding site (RBS), a 6-His tag and restriction enzyme sites.

The pQE vector was digested with the restriction enzymes indicated. The amplified sequences were ligated into the respective pQE vector and inserted in frame with the sequence

encoding for the RBS. The native stop codon was incorporated so the genes were not fused to the His tag of the vector. The ligation mixture was then used to transform the $E.\ coli$ Inc.) by electroporation. strain M15/pREP4 (Qiagen, M15/pREP4 contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants were identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies were selected. Plasmid DNA was isolated and confirmed by restriction analysis. Clones containing the desired constructs were grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture was used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells were grown to an optical density 600 (O.D.600) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalacto pyranoside") was then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression. Cells were grown an extra 3 to 4hours. Cells were then harvested by centrifugation.

The primer sequences set out above may also be employed to isolate the target gene from the deposited material by hybridization techniques described above.

Example 2

<u>Isolation of A Selected Clone From the Deposited Genomic</u> Clones

A clone is isolated directly by screening the deposited material using the oligonucleotide primers set forth in Example 1 for the particular gene desired to be isolated. The specific oligonucleotides are synthesized using an Applied Biosystems DNA synthesizer.

The two oligonucleotide primers corresponding to the gene of interest are used to amplify the gene from the deposited material. A polymerase chain reaction is carried out in 25 μ l of reaction mixture with 0.1 ug of the DNA of the gene of interest. The reaction mixture is 1.5-5 mM MgCl,, 0.01% (w/v) gelatin, 20 μM each of dATP, dCTP, dGTP,-dTTP, 25 pmol of each primer and 1.25 Unit of Taq polymerase. Thirty cycles of PCR (denaturation at 94°C for 1 min; annealing at 55°C for 1 min; elongation at 72°C for 1 min) are performed with the Perkin-Elmer Cetus 9600 thermal cycler. amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the gene of interest by subcloning and sequencing the DNA product. The ends of the newly purified genes are nucleotide sequenced to identify full length sequences. Complete sequencing of full length genes is then performed by Exonuclease III digestion or primer walking.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, within the scope of the appended claims, the invention may be practiced otherwise than as particularly described.

SEQUENCE LISTING

GENERAL INFORMATION: (1) (i) APPLICANT: RECOMBINANT BIOCATALYSIS, INC. TITLE OF INVENTION: (ii) THERMOSTABLE PHOSPHATASES (iii) NUMBER OF SEQUENCES: 54 CORRESPONDENCE ADDRESS: (iv) (A) ADDRESSEE: FISH & RICHARDSON (B) STREET: 4225 EXECUTIVE SQUARE, STE. 1400 (C) CITY: LA JOLLA (D) STATE: CA (E) COUNTRY: USA (F) ZIP: 92037 COMPUTER READABLE FORM: (v) (A) MEDIUM TYPE: 3.5 INCH DISKETTE (B) COMPUTER: IBM PS/2 (C) OPERATING SYSTEM: MS-DOS (D) SOFTWARE: WORD PERFECT 6.0 (vi) CURRENT APPLICATION DATA: (A) APPLICATION NUMBER: Unassigned (B) FILING DATE: June 19, 1997 (C) CLASSIFICATION: Unassigned (vii) PRIOR APPLICATION DATA: (A) APPLICATION NUMBER: (B) FILING DATE: (C) CLASSIFICATION: ATTORNEY/AGENT INFORMATION: (viii) (A) NAME: Haile, Lisa A. (B) REGISTRATION NUMBER: 38,347

(C) REFERENCE/DOCKET NUMBER: 09010/015W01

(ix)

TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 619-678-5070
(B) TELEFAX: 619-678-5099

(2)	INFORMATI	ON FOR SEQ ID NO:1:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 52 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:1:	
CCGAG.	AATTC ATTA	AAGAGG AGAAATTAAC TATGGGGGCA GGTCCGAAAA GG	52
(2)	INFORMATI	CON FOR SEQ ID NO:2:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 31 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:2:	
CCGAG	GATCC TCAG	CCGCCCC CTGCGGGTGC G	31
(2)	INFORMAT	ION FOR SEQ ID NO:3:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 52 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: CDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:3:	
CCGAG	BAATTC ATT.	AAAGAGG AGAAATTAAC TATGTTGGAT ATACTGCTTG TT	52
(2)	INFORMAT	ION FOR SEQ ID NO:4:	
	(i) -	SEQUENCE CHARACTERISTICS (A) LENGTH: 32 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:4:	
CCGA	GGATCC TTA	TTTTTTA ACCAAATTTC CC	32

(2)	INFORMATI	ON FOR SEQ ID NO:5:	-
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 52 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:5:	
CCGAC	AATTG ATTA	AAGAGG AGAAATTAAC TATGATGATG GAATTCACTC GC	52
(2)	INFORMATI	ON FOR SEQ ID NO:6:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 32 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:6:	
CGGAG	GATCC CTAC	CAGTTCT AAAAGTCTTT TA	32
(2)	INFORMATI	ON FOR SEQ ID NO:7:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 52 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:7:	
CCGAC	AATTG ATT	AAAGAGG AGAAATTAAC TATGAGAACC CTGACAATAA AC	52
(2)	INFORMAT	ION FOR SEQ ID NO:8:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 31 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:8:	

CCGAGGATCC TTACACCCAC AGAACCCTTA C

31

(2)	INFORMATI	ON FOR SEQ ID NO:9:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 52 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:9:	
CCGAG	AATTC ATT	AAAGAGG AGAAATTAAC TATGAAAGGA AAGTCTCTTG TT	52
(2)	INFORMAT:	ION FOR SEQ ID NO:10:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 31 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi) SEQ	UENCE DESCRIPTION: SEQ ID NO:10:	
CCGAG	GATCC TCA	AGCTTCC TGGAGAATCA A	31
(2)	INFORMAT	ION FOR SEQ ID NO:11:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 52 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: CDNA	
	(xi) SEQ	UENCE DESCRIPTION: SEQ ID NO:11:	
CCGAG	GAATTC ATT	AAAGAGG AGAAATTAAC TATGCCAAGA AATATCGCCG CT	52
(2)	INFORMAT	CION FOR SEQ ID NO:12:	
	(i) -	SEQUENCE CHARACTERISTICS (A) LENGTH: 34 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: CDNA	
	(xi) SEQ	QUENCE DESCRIPTION: SEQ ID NO:12:	
CGGA	GGATCC TT	AAGGCTTC TCGAGGTGGG GGTT	34

(2)	INFORMA	ATION FOR SEQ ID NO:13:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 52 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi) SI	EQUENCE DESCRIPTION: SEQ ID NO:13:	
CCGAC	AATTG A	TTAAAGAGG AGAAATTAAC TATGTATAAA TGGATTATTG AGGG	54
(2)	INFORM	ATION FOR SEQ ID NO:14:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 34 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: CDNA	
	(xi) S	EQUENCE DESCRIPTION: SEQ ID NO:14:	
CCGAG	GATCC C	TAAACATAG TCTAAGTAAT TAGC	34
(2)	INFORM	ATION FOR SEQ ID NO:15:	
	(i)	SEQUENCE CHARACTERISTICS (A) LENGTH: 52 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: CDNA	
	(xi) S	EQUENCE DESCRIPTION: SEQ ID NO:15:	
CCGAC	BAATTC A	TTAAAGAGG AGAAATTAAC TATGAGAATC CTCCTCACCA AC	32
(2)	INFORM	NATION FOR SEQ ID NO:16:	
-	(i) -	SEQUENCE CHARACTERISTICS (A) LENGTH: 31 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR	
	(ii)	MOLECULE TYPE: cDNA	
	(xi) S	SEQUENCE DESCRIPTION: SEQ ID NO:16:	
CCGA	GGATCC T	CACAGGCTC AGAAGCCTTT G	31

(2)	INFORM	IATION F	OR SE	aı Ç	NO:	17:									
	(i)	(A) (B) (C)	UENCE LENG TYPE STRAI TOPO	TH: : N NDED	54 UCLE NESS	NUCL IC A : S	EOTI CID INGL	DES							
	(ii)	MOL	ECULE	TYP	E: '	GENO	MIC :	DNA							
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:														
CCGAGA	CCGAGAATTC ATTAAAGAGG AGAAATTAAC TATGGAAAAC TTAAAAAAAGT ACCT														
(2)	2) INFORMATION FOR SEQ ID NO:18:														
	(i) SEQUENCE CHARACTERISTICS (A) LENGTH: 31 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR														
	(ii)	MOL	ECULE	TYP	E :	CDNA									
	(xi)	SEQ	UENCE	DES	CRIP	TION	: S	EQ I	D NO	:18:					
CGGAA	CGGAAGATCT TCACACCGCC ACTTCCATAT A														
(2)	INFORM	ATION E	OR SE	EQ II	NO:	19:									
	(i)	SEC	UENCE	CHA	RACT	ERIS	TICS								
	(i) SEQUENCE CHARACTERISTICS (A) LENGTH: 783 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE (D) TOPOLOGY: LINEAR														
	(ii)	MOL	ECULE	TYP	Ε:	geno	mic	DNA							
	(xi)	SEC	UENCE	DES	CRIP	TION	: s	EQ I	D NO	:19:					
ATG A	GG GGG	AGC GG	A GTG	CGG	ATA	CTT	CTC	ACC	AAC	GAT	GAC	GGC	ATC	48	
TTT G	CC GAG	GGT CT	GGG	GCT	CTG	CGC	AAG	ATG	CTG	GAG	CCC	GTG	GCT	96	
ACC C	TT TAC	GTG GT	GCT	CCG	GAC	CGA	GAG	CGT	AGC	GCG	GCC	AGC	CAT	144	
GCT A	TC ACC	GTT CA	c ccc	CCC	CTG	CGG	GTG	CGG	GAG	GCG	GGT	TTT	CGC	192	
AGC C	CC AGG	CTT AA	A GGC	TGG	GTA	GTG	GAC	GGT	ACC	CCG	GCC	GAC	TGC	240	
GTC A	ag ctg	GGC CT	g gag	GTA	CTT	TTG	CCC	GAA	CGT	CCA	GAT	TTC	CTG	288	
GTT T	CG GGC	ATA AA	C TAC	GGG	CCC	AAC	CTG	GGT	ACC	GAC	GTA	CTT	TAC	336	
TCC G	GC ACC	GTC TC	G GCG	GCC	ATA	GAA	GGG	GTA	ATT	AAC	GGC	ATT	ccc	384	
TCG G	TG GCC	GTA TC	r TTG	GCC	ACG	CGG	CGG	GAG	CCG	GAC	TAT	ACC	TGG	432	
GCG G	CC CGG	TTC GT	CTG	GTC	CTG	CTG	GAG	GAA	CTG	CGA	AAA	CAC	CAA	480	

CTG CCC CCA GGA ACC CTG CTC AAC GTC AAC GTG CCC GAC GGG GTG CCC 528

CGC GGG GTC AAG GTG ACC AAA CTG GGA AGC GTA CGC TAC GTC AAC GTG 576GTA GAC TGC CGC ACC GAC CCT CGG GGG AAG GCT TAC TAC TGG ATG GCG 624

GGA GAA CCA TTG GAG CTG GAC GGC AAC GAC TCC GAA ACC GAC GTC TGG 672

GCG GTG CGA GAA GGC TAT ATT TCC GTA ACA CCG GTC CAG ATC GAC CTT 720

ACT AAC TAC GGC TTC CTG GAA GAA CTC AAA AAA TGG CGT TTC AAG GAT 768

ATC TTT TCT TCT TAA

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 765 NUCLEOTIDES
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: genomic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATG TTG GAT ATA CTG CTT GTT AAT GAT GGC ATT TAT TCA AAT GGA 48 TTA ATA GCT TTG AAG GAT GCA TTA TTG GAA AAA TTT AAT GCG AGG ATT 96 ACT ATT GTA GCC CCA ACA AAT CAG CAG AGT GGT ATT GGT AGG GCA ATA 144 AGT TTA TTC GAG CCG TTA AGG ATA ACT AAA ACC AAA TTA GCA GAT GGT 192 TCT TGG GGA TAT GCA GTT TCA GGA ACC CCA ACA GAT TGC GTT ATA TTG 240 GGC ATT TAT GAG ATA TTA AAG AAG GTA CCT GAT GTA GTT ATA TCA GGA 288 ATA AAC ATT GGA GAA AAC CTT GGG ACT GAA ATA ACA ACT TCT GGA ACG 336 TTG GGG GCT GCG TTT GAA GGG GCC CAT CAT GGG GCT AAG GCA TTA GCA 384 TCA TCA CTC CAA GTT ACC TCT GAC CAT CTA AAG TTT AAA GAG GGG GAG 432 ACC CCA ATA GAC TTC ACA GTC CCA GCA AGA ATT ACT GCA AAT GTT GTT 480 GAG AAG ATG TTG GAT TAT GAT TTC CCA TGT GAT GTC GTC AAC TTA AAC 528 ATT CCA GAA GGA GCA ACA GAA AAG ACA CCG ATT GAA ATC ACA AGG TTG 576 GCA AGG AAA ATG TAT ACA ACA CAC GTT GAG GAA AGA ATA GAT CCA AGA 624 GGG AGG AGT TAT TAT TGG ATT GAT GGG TAT CCT ATT TTA GAG GAA GAG 672 GAA GAC ACT GAT GTC TAT GTT GTT AGA AGA AAG GGA CAT ATT TCT CTA 720 ACC CCA TTA ACA TTA GAC ACA ACA ATT AAA AAT TTA GAG GAA TTT AAG 768 798 AAA AAA TAT GAG AGA ATA TTA AAT GAA TGA

INFORMATION FOR SEQ ID NO:21: (2)

- SEQUENCE CHARACTERISTICS (i) (A) LENGTH: 765 NUCLEOTIDES (B) TYPE: NUCLEIC ACID (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR
- (ii)MOLECULE TYPE: genomic DNA
- SEQUENCE DESCRIPTION: SEQ ID NO:21: (xi)

ATG ATG ATG GAA TTC ACT CGC GAG GGA ATA AAA GCT GCT GTA GAG GCA 48 CTT CAA GGG TTA GGA GAG ATC TAC GTA GTT GCC CCA ATG TTT CAA AGG 96 AGC GCA AGT GGA AGG GCA ATG ACC ATC CAC AGA CCT CTA AGG GCT AAA 144 AGA ATA AGT ATG AAC GGT GCA AAA GCA GCC TAT GCT TTG GAT GGA ATG 192 CCC GTT GAT TGC GTT ATC TTT GCC ATG GCC AGA TTT GGA GAT TTC GAC 240 CTT GCA ATA AGT GGT GTA AAC TTG GGA GAA AAC ATG AGC ACC GAG ATA 288 ACG GTT TCC GGG ACT GCA AGC GCT GCA ATA GAG GCT GCA ACC CAA GAG 336 ATC CCA AGC ATT CCC ATA AGC CTG GAA GTT AAT AGA GAA AAA CAC AAA 384 TTT GGT GAG GGC GAA GAG ATT GAC TTC TCA GCT GCC AAG TAT TTC CTA 432 AGA AAA ATC GCA ACG GCG GTT TTA AAG AGA GGC CTC CCC AAA GGA GTC 480 GAT ATG CTG AAC GTC AAC GTC CCT TAT GAT GCA AAT GAA AGG ACA GAG 528 ATA GCT TTT ACT CGC CTG GCA AGA AGG ATG TAT AGG CCT TCT ATT GAA 576 GAG CGC ATA GAC CCA AAG GGG AAT CCC TAC TAC TGG ATA GTT GGA ACT 624 CAG TGC CCT AAG GAG GCA TTA GAG CCG GGA ACG GAT ATG TAT GTA GTT 672 AAA GTT GAG AGA AAA GTT AGC GTG ACT CCA ATA AAC ATT GAT ATG ACA 720 GCA AGA GTG AAT TTA GAC GAG ATT AAA AGA CTT TTA GAA CTG TAG 765

INFORMATION FOR SEQ ID NO:22: (2)

- SEQUENCE CHARACTERISTICS (i)
 - (A) LENGTH: 816 NUCLEOTIDES (B) TYPE: NUCLEIC ACID

 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR
- MOLECULE TYPE: genomic DNA (ii)
- SEQUENCE DESCRIPTION: SEQ ID NO:22: (xi)

ATG AGA ACC CTG ACA ATA AAC ACT GAC GCG GAG GGG TTC GTT TTG AGG

48

ATT CTC CTG ACG AAC GAC GAT GGA ATC TAC TCC AAC GGA CTG CGC GCC 96-GCT GTG AAA GCC CTG AGT GAG CTC GGC GAA GTT TAC GTC GTT GCC CCC 144 CTC TTC CAG AGG AGC GCG AGC GGC AGG GCC ATG ACG CTC CAC AGG CCG 192 ATA AGG GCC AAG CGC GTT GAC GTT CCC GGC GCA AAG ATA GCC TAC GGA 240 ATA GAT GGA ACT CCT ACT GAC TGC GTG ATT TTC GCC ATA GCC CGC TTC 288 GGG AGC TTT GGT TTA GCC GTG AGC GGG ATT AAC CTC GGC GAG AAC CTG 336 AGC ACC GAG ATA ACA GTC TCA GGG ACG GCC TCC GCT GCC ATA GAG GCC 384 TCA ACT CAT GGA ATT CCG AGC ATA GCG ATT AGC CTT GAG GTG GAG TGG 432 AAG AAG ACC CTC GGC GAG GGT GAG GGG GTT GAC TTC TCG GTC TCG ACT 480 CAC TTC CTC AAG AGA ATC GCG GGA GCC CTC TTG GAG AGA GGT CTT CCT 528 GAG GGC GTT GAC ATG CTC AAC GTC AAC GTT CCG AGC GAC GCG ACG GAG 576 GAA ACG GAG ATA GCA ATC ACC CGC TTA GCC CGG AAG CGC TAC TCC CCA 624 ACG GTC GAG GAG AGG ATT GAC CCC AAG GGC AAC CCC TAC TAC TGG ATT 672 GTC GGC AAA CTT GTC CAA GAC TTC GAG CCA GGG ACA GAT GCC TAC GCC 720 CTG AAG GTC GAG AGG AAG GTC AGC GTC ACG CCG ATA AAC ATA GAT ATG 768 ACT GCG AGG GTG GAC TTT GAG GAG CTT GTA AGG GTT CTG TGG GTG TAA 816

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 1494 NUCLEOTIDES
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: genomic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

ATG AAA GGA AAG TCT CTT GTT AGC GGT CTG TTG TTG GGT CTT TTA ATT 48 TTG AGC CTG ATT TCA TTC CAG CCA AGC TTT GCA TAC TCC CCA CAC GGC 96 GGT GTC AAA AAC ATC ATA ATC CTG GTT GGA GAC GGC ATG GGT CTT GGG 144 CAT GTA GAA ATT ACA AAG CTC GTT TAT GGA CAC TTA AAC ATG GAA AAC 192 TTT CCA GTT ACT GGA TTT GAG CTT ACT GAT TCC CTA AGT GGT GAA GTT 240 ACA GAT TCT GCT GCG GCA GGA ACT GCA ATA TCC ACT GGA GCT AAA ACG 288 TAT AAT GGT ATG ATT TCA GTA ACC AAC ATA ACC GGA AAG ATA GTT AAC 336 TTA ACA ACC CTA CTT GAA GTG GCT CAA GAG CTT GGG AAG TCA ACA GGG 384 CTG GTC ACC ACA ACA AGG ATT ACC CAT GCA ACT CCA GCA GTT TTT GCG 432

PCT/US97/10784 WO 97/48416

TCC	CAT	GTC	CCA	GAT	AGG	GAT	ATG	GAG	GGG	GAG	ATA	CCC	AAG	CAA	CTC	480-
ATA	ATG	CAC	AAA	GTT	AAC	GTC	TTG	TTG	GGT	GGT	GGA	AGG	GAG	AAA	TTC	528
GAT	GAG	AAA	AAT	TTG	GAG	CTG	GCC	AAA	AAG	CAG	GGA	TAC	AAA	GTA	GTT	576
TTC	ACG	AAG	GAA	GAG	CTT	GAA	AAA	GTT	GAA	GGA	GAT	TAT	GTC	CTA	GGA	624
CTC	TTT	GCA	GAA	AGT	CAC	ATC	CCT	TAC	GTA	TTG	GAT	AGA	AAA	CCC	GAT	672
GAT	GTT	GGA	CTT	ATT	GAA	ATG	GCC	AAA	AAG	GCA	ATT	TCA	ATA	CTC	GAG	720
AAG	AAC	CCG	AGC	GGA	TTC	TTT	CTC	ATG	GTT	GAG	GGC	GGA	AGG	ATT	GAC	768
CAT	GCA	GCC	CAT	GGA	AAC	GAT	GTC	GCA	TCG	GTT	GTT	GCA	GAA	ACT	AAG	816
GAG	TTT	GAC	GAT	GTT	GTC	AGA	TAC	GTG	CTG	GAA	TAT	CCG	AAG	AAG	AGG	864
GGA	GAT	ACC	TTG	GTA	ATA	GTG	CTT	GCC	GAT	CAC	GAA	ACT	GGA	GGT	CTT	912
GCA	ATA	GGT	CTA	ACG	TAT	GGA	AAT	GCA	ATC	GAT	GAA	GAT	GCC	ATA	AGA	960
AAA	ATA	AAA	GCA	AGC	ACG	TTG	AGG	ATG	CCC	AAA	GAG	GTT	AAG	GCA	GGG	1008
AGT	AGT	GTA	AAA	GAG	TCC	TCA	AAG	GTA	TGC	CGG	ATT	TGT	CCC	AAC	AGA	1056
GGA	AGA	AGT	CAG	TAT	ATT	GAG	AAT	GCG	CTG	CAC	TCG	ACA	AAC	AAG	TAT	1104
GCC	CTC	TCA	AAT	GCA	GTA	GCC	GAT	GTT	ATA	AAC	AGG	CGT	ATT	GGT	GTT	1152
GGA	TTC	ACC	TCC	TAT	GAG	CAT	ACA	GGA	GTT	CCA	GTT	CCG	CTC	TTA	GCT	1200
TAC	GGT	CCC	GGG	GCA	GAG	AAC	TTC	AGA	GGT	TTC	TTA	CAC	CAT	GTG	GAT	1248
ACA	GCA	AGA	TTA	GTT	GCA	AAG	TTA	ATG	CTC	TTT	GGA	AGG	AGG	AAT	ATT	1296
CCA	GTT	ACC	ATT	TCA	AGC	GTG	AGC	AGT	GTT	AAG	GGA	GAC	ATA	ACC	GGT	1344
GAT	TAC	AGG	GTT	GAT	GAG	AAG	GAT	GCC	TAC	GTT	ACG	CTC	ATG	ATG	TTT	1392
CTC	GGA	GAA	AAA	GTG	GAT	AAT	GAA	ATT	GAA	AAG	AGA	GTC	GAT	ATA	GAC	1440
AAC	AAC	GGC	ATG	GTT	GAC	TTA	AAT	GAC	GTC	ATG	TTG	ATT	CTC	CAG	GAA	1488
GCT	TGA															1494

INFORMATION FOR SEQ ID NO:24: . (2)

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 1755 NUCLEOTIDES

(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: SINGLE

(D) TOPOLOGY: LINEAR

MOLECULE TYPE: genomic DNA (ii)

SEQUENCE DESCRIPTION: SEQ ID NO:24: (xi)

ATG CCA AGA AAT ATC GCC GCT GTA TGC GCC CTG GCC GCT TTG TTA GGG 48 TCG GCC TGG GCG GCC AAA GTT GCC GTC TAC CCC TAC GAC GGA GCC GCT 96 TTG CTG GCG GGG CAG CGC TTC GAT TTG CGC ATA GAA GCC TCC GAG CTG 144

AAA	GGC	AAT	TTA	AAG	GCT	TAC	CGC	ATC	ACC	CTG	GAC	GGC	CAG	CCT	CTG	192 -
GCG	GGC	CTC	GAG	CAA	ACC	GCG	CAG	GGG	GCC	GGG	CAG	GCC	GAG	TGG	ACC	240
CTG	CGC	GGT	GCC	TTC	CTG	CGC	CCT	GGA	AGC	CAC	ACC	CTC	GAG	GTC	AGC	288
CTC	ACC	GAC	GAC	GCT	GGG	GAG	AGC	AGG	AAG	AGC	GTA	CGT	TGG	GAG	GCT	336
CGG	CAG	AAC	CTT	CGC	TTG	CCC	CGA	GCG	GCC	AAG	AAT	GTG	ATT	CTC	TTC	384
ATT	GGC	GAC	GGG	ATG	GGC	TGG	AAC	ACC	CTC	AAC	GCC	GCC	CGC	ATC	ATC	432
GCC	AAA	GGC	TTT	AAC	CCC	GAA	AAC	GGT	ATG	ccc	AAC	GGA	AAC	CTC	GAG	480
ATC	GAG	AGT	GGT	TAC	GGT	GGG	ATG	GCT	ACC	GTC	ACT	ACC	GGC	AGC	TTT	528
GAT	AGC	TTC	ATC	GCC	GAC	TCA	GCT	AAC	TCG	GCT	TCT	TCC	ATC	ATG	ACC	576
GGG	CAG	AAG	GTG	CAG	GTG	AAT	GCC	CTC	AAC	GTT	TAC	CCA	TCA	AAC	CTC	624
AAA	GAT	ACC	CTG	GCC	TAC	CCC	CGG	ATC	GAA	ACC	CTA	GCG	GAG	ATG	CTC	672
AAG	CGG	GTA	CGC	GGG	GCC	AGC	ATT	GGG	GTA	GTG	ACC	ACC	ACC	TTC	GGC	720
ACC	GAC	GCT	ACC	CCG	GCT	TCA	CTC	AAC	GCC	CAT	ACC	CGC	CGC	CGC	GGT	768
GAT	TAC	CAG	GCT	ATC	GCC	GAC	ATG	TAC	TTT	GGT	AGA	GGC	GGG	TTC	GGT	816
GTT	CCC	TTG	GAT	GTG	ATG	CTC	TTC	GGT	GGT	TCA	CGC	GAC	TTC	ATC	ccc	864
CAG	AGC	ACC	CCT	GGC	TCG	CGG	CGC	AAG	GAT	AGC	ACG	GAC	TGG	ATT	GCC	912
GAA	TCC	CAG	AAG	CTG	GGC	TAC	ACC	TTT	GTC	AGC	ACC	CGC	AGC	GAG	CTG	960
CTG	GCG	GCC	AAA	CCC	ACC	GAT	AAG	CTG	TTT	GGG	CTG	TTC	AAC	ATT	GAC	1008
AAC	TTC	CCC	AGC	TAC	CTA	GAC	CGC	GCA	GTG	TGG	AAG	CGG	CCC	GAG	ATG	1056
CTG	GGA	AGC	TTT	ACC	GAT	ATG	CCC	TAC	CTC	TGG	GAG	ATG	ACC	CAG	AAA	1104
GCC	GTG	GAG	GCT	CTC	TCC	AGA	AAC	GAC	AAA	GGC	TTT	TTC	TTG	ATG	GTT	1152
GAG	GGG	GGA	ATG	GTG	GAT	AAG	TAC	GAG	CAC	CCC	TTG	GAC	TGG	CCC	CGC	1200
GCA	CTT	TGG	GAT	GTA	CTC	GAG	CTG	GAC	CGC	GCG	GTG	GCT	TGG	GCC	AAG	1248
GGC	TAT	GCG	GCC	TCC	CAC	CCC	GAT	ACC	CTG	GTG	ATT	GTC	ACC	GCC	GAC	1296
CAC	GCT	CAC	TCG	ATC	TCG	GTG	TTT	GGC	GGT	TAC	GAC	TAC	TCC	AAG	CAG	1344
GGC	ČGG	GAG	GGG	GTG	GGG	GTT	TAT	GAG	GCC	GCC	AAG	TTC	CCC	ACC	TAC	1392
GGC	GAC	AAA	AAA	GAC	GCC	AAC	GGC	TTT	CCC	TTG	CCC	GAC	ACC	ACT	CGG	1440
GGA	ATC	GCG	GTA	GGC	TTC	GGG	GCC	ACG	CCG	GAT	TAC	TGT	GAA	ACC	TAC	1488
CGG	GGC	CGC	GAG	GTC	TAC	AAA	GAC	CCC	ACC	ATC	TCC	GAC	GGC	AAA	GGT	1536
GGT	TAC	GTG	GCC	AAC	CCT	GAG	GTC	TGC	AAG	GAG	CCG	GGC	CTT	CCA	ACG	1584
TAC	CGG	CAA	CTC	CCA	GTA	GAT	AGC	GCC	CAG	GGC	GTG	CAC	ACG	GCT	GAT	1632
CCC	ATG	CCG	CTG	TTT	GCC	TTT	GGC	GTG	GGG	TCT	CAG	TTC	TTC	AAT	GGC	1680

CTC ATC GAC CAG ACC GAG ATC TTC TTC CGC ATG GCC CAG GCC CTA GGG 1728TTC AAC CCC CAC CTC GAG AAG CCT TAA 1755

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS
 (A) LENGTH: 912 NUCLEOTIDES
 - (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: SINGLE
 - (C) STRANDEDNESS: SINGL
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: genomic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

ATG TAT AAA TGG ATT ATT GAG GGT AAG CTT GCC CAA GCA CCT TTT CCA 48 AGC CTA GGT GAA CTA GCC GAT CTC AAA AGA CTT TTC GAC GCC ATT ATT 96 GTT CTT ACA ATG CCG CAT GAA CAA CCG CTT AAT GAG AAA TAT ATC GAG 144 ATA TTA GAG AGC CAT GGA TTC CAA GTC CTC CAT GTC CCC ACG CTC GAC 192 TTT CAT CCT TTA GAA CTC TTC GAC CTT TTG AAA ACA AGC ATA TTC ATT 240 GAT GAA AAC CTG GAG AGA TCC CAC AGA GTG CTT GTC CAC TGC ATG GGA 288 GGC ATA GGC CGG AGC GGG CTT GTA ACT GCT GCG TAC TTA ATA TTC AAA 336 GGT TAT GAT ATT TAC GAC GCG GTA AAG CAT GTG AGA ACG GTA GTG CCT 384 GGT GCT ATT GAA AAC AGA GGG CAA GCG TTA ATG CTT GAG AAC TAC TAT 432 ACC CTG GTC AAA AGT TTC AAC AGA GAG TTG CTG AGA GAC TAC GGG AAG 480 AAA ATT TTC ACG CTC GGT GAC CCG AAG GCG GTT CTC CAC GCT TCT AAG 528 ACG ACT CAG TTC ACG ATT GAA CTC TTA AGC AAC TTA CAC GTC AAC GAG 576 GCG TTT TCA ATC AGT GCG ATG GCT CAA TCA CTG CTC CAC TTT CAC GAC 624 GTA AAA GTC CGC TCT AAA CTG AAA GAA GTA TTC GAA AAC ATG GAA TTC 672 TCA TCC GCC TCA GAG GAG GTT CTG TCA TTT ATT CAC CTA CTC GAT TTC 720 TAT CAG GAT GGC AGG GTT GTT TTA ACC ATT TAC GAT TAT CTC CCC GAT 768 AGG GTG GAT TTG ATT TTA TTG TGT AAG TGG GGT TGT GAT AAA ATA GTT 816 GAA GTC TCG TCT TCA GCG AAG AAA ACC GTT GAG AAG CTT GTA GGA AGA 864 AAG GTT TCC CTA TCC TGG GCT AAT TAC TTA GAC TAT GTT TAG 912

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 774 NUCLEOTIDES
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

PCT/US97/10784 WO 97/48416

'ii)	MOLECULE	TVDE.	genomic	DNA
1 1 1	PICHELONE	LIPEL	d CIIOniz c	

	(x	i)		SEQU	ENCE	DES	CRIP	TION	: S	EQ I	D NO	:26:				
ATG	AGA	ATC	CTC	CTC	ACC	AAC	GAC	GAC	GGC	ATC	TAT	TCC	AAC	GGT	CTG	48
CGC	GCG	GCG	GTG	AAG	GGC	CTG	AGC	GAG	CTC	GGC	GAG	GTC	TAC	GTC	GTC	96
GCC	CCG	CTC	TTC	CAG	AGG	AGC	GCG	AGC	GGT	CGG	GCG	ATG	ACC	CTA	CAC	144
AGG	CCG	ATA	AGG	GCA	AAG	AGG	GTT	GAC	GTT	CCC	GGC	GCG	AAG	ATA	GCG	192
TAT	GGC	ATA	GAC	GGA	ACG	CCG	ACC	GAC	TGC	GTG	ATT	TTT	GCC	ATC	GCC	240
CGC	TTC	GGC	GAC	TTT	GAT	CTG	GCG	GTC	AGC	GGG	ATA	AAC	CTA	GGC	GAG	288
AAC	CTG	AGC	ACG	GAG	ATA	ACC	GTC	TCC	GGA	ACG	GCC	TCG	GCG	GCG	ATA	336
GAG	GCT	TCC	ACC	CAC	GGG	ATT	CCA	AGT	GTA	GCT	ATA	AGC	CTC	GAG	GTC	384
GAG	TGG	AAG	AAG	ACC	CTC	GGC	GAG	GGG	GAG	GGT	ATT	GAC	TTC	TCG	GTT	432
TCA	GCA	CAC	TTC	CTG	AGA	AGG	ATA	GCG	ACG	GCT	GTC	CTT	AAG	AAG	GGC	480
CTG	CCT	GAA	GGG	GTG	GAC	ATG	CTC	AAC	GTG	AAC	GTC	CCT	AGC	GAC	GCC	528
AGC	GAG	GGG	ACT	GAG	ATC	GCC	ATA	ACG	CGC	CTC	GCG	AGG	AAG	CGC	TAT	576
TCT	CCG	ACG	ATA	GAG	GAG	AGG	ATA	GAC	CCC	AAG	GGC	AAC	CCC	TAC	TAC	624
TGG	ATC	GTT	GGC	AGG	CTC	GTC	CAG	GAG	TTC	GAG	CCG	GGC	ACG	GAC	GCC	672
TAC	GCT	CTG	AAA	GTC	GAG	AGA	AAG	GTC	AGC	GTC	ACG	CCC	ATA	AAC	ATC	720
GAC	ATG	ACT	GCG	AGG	GTT	GAC	TTT	GAG	AAC	CTT	CAA	AGG	CTT	CTG	AGC	768
CTG	TGA															774

INFORMATION FOR SEQ ID NO:27: (2)

- SEQUENCE CHARACTERISTICS (i)
 - (A) LENGTH: 795 NUCLEOTIDES (B) TYPE: NUCLEIC ACID

 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR
- MOLECULE TYPE: genomic DNA (ii)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

ATG GAA AAC TTA AAA AAG TAC CTA GAA GTT GCA AAA ATA GCC GCG CTC 48 GCG GGT GGG CAG GTT CTG AAA GAA AAC TTC GGA AAG GTA AAA AAG GAA 96 AAC ATA GAG GAA AAA GGG GAA AAG GAC TTT GTA AGT TAC GTG GAT AAA 144 ACT TCA GAG GAA AGG ATA AAG GAG GTG ATA CTC AAG TTC TTT CCC GAT 192 CAC GAG GTC GTA GGG GAA GAG ATG GGT GCG GAG GGA AGC GGA AGC GAA 240 TAC AGG TGG TTC ATA GAC CCC CTT GAC GGC ACA AAG AAC TAC ATA AAC 288

PCT/US97/10784 WO 97/48416

GGT	TTT	CCC	ATC	TTT	GCC	GTA	TCA	GTG	GGA	CTT	GTT	AAG	GGA	GAA	GAG	336
CCA	ATT	GTG	GGT	GCG	GTT	TAC	CTT	CCT	TAC	TTT	GAC	AAG	CTT	TAC	TGG	384
GGT	GCT	AAA	GGT	CTC	GGG	GCT	TAC	GTA	AAC	GGA	AAG	AGG	ATA	AAG	GTA	432
AAG	GAC	AAT	GAG	AGT	TTA	AAG	CAC	GCC	GGA	GTG	GTT	TAC	GGA	TTT	CCC	480
TCT	AGG	AGC	AGG	AGG	GAC	ATA	TCT	ATC	TAC	TTG	AAC	ATA	TTC	AAG	GAT	528
GTC	TTT	TAC	GAA	GTT	GGC	TCT	ATG	AGG	AGA	CCC	GGG	GCT	GCT	GCG	GTT	576
GAC	CTC	TGC	ATG	GTG	GCG	GAA	GGG	ATA	TTT	GAC	GGG	ATG	ATG	GAG	TTT	624
GAA	ATG	AAG	CCG	TGG	GAC	ATA	ACC	GCA	GGG	CTT	GTA	ATA	CTG	AAG	GAA	672
GCC	GGG	GGC	GTT	TAC	ACA	CTT	GTG	GGA	GAA	CCC	TTC	GGA	GTT	TCG	GAC	720
ATA	ATT	GCG	GGC	AAC	AAA	GCC	CTC	CAC	GAC	TTT	ATA	CTT	CAG	GTA	GCC	768
AAA	AAG	TAT	ATG	GAA	GTG	GCG	GTG	TGA								799

INFORMATION FOR SEQ ID NO:28: (2)

- SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 260 AMINO ACIDS

 - (B) TYPE: AMINO ACID (D) TOPOLOGY: LINEAR
- (ii)MOLECULE TYPE: PROTEIN
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Met Arg Gly Ser Gly Val Arg Ile Leu Leu Thr Asn Asp Asp Gly Ile

Phe Ala Glu Gly Leu Gly Ala Leu Arg Lys Met Leu Glu Pro Val Ala 20 25 30

Thr Leu Tyr Val Val Ala Pro Asp Arg Glu Arg Ser Ala Ala Ser His 35 40 45

Ala Ile Thr Val His Arg Pro Leu Arg Val Arg Glu Ala Gly Phe Arg

Ser Pro Arg Leu Lys Gly Trp Val Val Asp Gly Thr Pro Ala Asp Cys 65 70 75 80

Val Lys Leu Gly Leu Glu Val Leu Leu Pro Glu Arg Pro Asp Phe Leu

Val Ser Gly Ile Asn Tyr Gly Pro Asn Leu Gly Thr Asp Val Leu Tyr

Ser Gly Thr Val Ser Ala Ala Ile Glu Gly Val Ile Asn Gly Ile Pro

Ser Val Ala Val Ser Leu Ala Thr Arg Arg Glu Pro Asp Tyr Thr Trp

Ala Ala Arg Phe Val Leu Val Leu Leu Glu Glu Leu Arg Lys His Gln 155

Leu Pro Pro Gly Thr Leu Leu Asn Val Asn Val Pro Asp Gly Val Pro

Ala Val Arg Glu Gly Tyr Ile Ser Val Thr Pro Val Gln Ile Asp Leu

Thr Asn Tyr Gly Phe Leu Glu Glu Leu Lys Lys Trp Arg Phe Lys Asp 245 250 255

Ile Phe Ser Ser

- (2) INFORMATION FOR SEQ ID NO:29:
 - (i) SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 265 AMINO ACIDS
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
 - (ii) MOLECULE TYPE: PROTEIN
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Met Leu Asp Ile Leu Leu Val Asn Asp Asp Gly Ile Tyr Ser Asn Gly 5 10 15

Leu Ile Ala Leu Lys Asp Ala Leu Leu Glu Lys Phe Asn Ala Arg Ile 20 25 30

Thr Ile Val Ala Pro Thr Asn Gln Gln Ser Gly Ile Gly Arg Ala Ile 35 40 45

Ser Leu Phe Glu Pro Leu Arg Ile Thr Lys Thr Lys Leu Ala Asp Gly 50 55 60

Ser Trp Gly Tyr Ala Val Ser Gly Thr Pro Thr Asp Cys Val Ile Leu 65 70 75 80

Gly Ile Tyr Glu Ile Leu Lys Lys Val Pro Asp Val Val Ile Ser Gly
- 85 90 95

Ile Asn Ile Gly Glu Asn Leu Gly Thr Glu Ile Thr Thr Ser Gly Thr 100 105 110

Leu Gly Ala Ala Phe Glu Gly Ala His His Gly Ala Lys Ala Leu Ala 115 120 125

Ser Ser Leu Gln Val Thr Ser Asp His Leu Lys Phe Lys Glu Gly Glu 130 140

Thr Pro Ile Asp Phe Thr Val Pro Ala Arg Ile Thr Ala Asn Val Val 145 150 155 160

Glu Lys Met Leu Asp Tyr Asp Phe Pro Cys Asp Val Val Asn Leu Asn 165 170 175

Ile Pro Glu Gly Ala Thr Glu Lys Thr Pro Ile Glu Ile Thr Arg Leu 180 185 190

Ala Arg Lys Met Tyr Thr Thr His Val Glu Glu Arg Ile Asp Pro Arg 195 200 205

Gly Arg Ser Tyr Tyr Trp Ile Asp Gly Tyr Pro Ile Leu Glu Glu Glu 210 220

Glu Asp Thr Asp Val Tyr Val Val Arg Arg Lys Gly His Ile Ser Leu 225 230 235

Thr Pro Leu Thr Leu Asp Thr Thr Ile Lys Asn Leu Glu Glu Phe Lys 245 250 255

Lys Lys Tyr Glu Arg Ile Leu Asn Glu 260 265

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 254 AMINO ACIDS
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Met Met Met Glu Phe Thr Arg Glu Gly Ile Lys Ala Ala Val Glu Ala

Leu Gln Gly Leu Gly Glu Ile Tyr Val Val Ala Pro Met Phe Gln Arg 20 25 30

Ser Ala Ser Gly Arg Ala Met Thr Ile His Arg Pro Leu Arg Ala Lys

Arg Ile Ser Met Asn Gly Ala Lys Ala Ala Tyr Ala Leu Asp Gly Met 50 60

Pro Val Asp Cys Val Ile Phe Ala Met Ala Arg Phe Gly Asp Phe Asp 65 70 75 80

Leu Ala Ile Ser Gly Val Asn Leu Gly Glu Asn Met Ser Thr Glu Ile 85 90 95

Thr Val Ser Gly Thr Ala Ser Ala Ala Ile Glu Ala Ala Thr Gln Glu 100 105 110

Ile Pro Ser Ile Pro Ile Ser Leu Glu Val Asn Arg Glu Lys His Lys 115 120 125

Phe Gly Glu Glu Glu Ile Asp Phe Ser Ala Ala Lys Tyr Phe Leu 130 135 140

Arg Lys Ile Ala Thr Ala Val Leu Lys Arg Gly Leu Pro Lys Gly Val 145 150 155 160

Asp Met Leu Asn Val Asn Val Pro Tyr Asp Ala Asn Glu Arg Thr Glu 165 170 175

Ile Ala Phe Thr Arg Leu Ala Arg Arg Met Tyr Arg Pro Ser Ile Glu 180 185 190

Glu Arg Ile Asp Pro Lys Gly Asn Pro Tyr Tyr Trp Ile Val Gly Thr 195 200 205

Gln Cys Pro Lys Glu Ala Leu Glu Pro Gly Thr Asp Met Tyr Val Val 210 215 220

Lys Val Glu Arg Lys Val Ser Val Thr Pro Ile Asn Ile Asp Met Thr 225 230 235 240

Ala Arg Val Asn Leu Asp Glu Ile Lys Arg Leu Leu Glu Leu
245

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 271 AMINO ACIDS
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Met Arg Thr Leu Thr Ile Asn Thr Asp Ala Glu Gly Phe Val Leu Arg
5 10 15

Ile Leu Leu Thr Asn Asp Asp Gly Ile Tyr Ser Asn Gly Leu Arg Ala
20 25 30

Ala Val Lys Ala Leu Ser Glu Leu Gly Glu Val Tyr Val Val Ala Pro 35 40 45

Leu Phe Gln Arg Ser Ala Ser Gly Arg Ala Met Thr Leu His Arg Pro 50 55 60

Ile Arg Ala Lys Arg Val Asp Val Pro Gly Ala Lys Ile Ala Tyr Gly 65 70 75 80

Ile Asp Gly Thr Pro Thr Asp Cys Val Ile Phe Ala Ile Ala Arg Phe 85 90 95

Gly Ser Phe Gly Leu Ala Val Ser Gly Ile Asn Leu Gly Glu Asn Leu
100 105 110

Ser Thr Glu Ile Thr Val Ser Gly Thr Ala Ser Ala Ala Ile Glu Ala 115 120 125

Ser Thr His Gly Ile Pro Ser Ile Ala Ile Ser Leu Glu Val Glu Trp 130 135 140

Lys Lys Thr Leu Gly Glu Gly Glu Gly Val Asp Phe Ser Val Ser Thr 145 150 155 160

His Phe Leu Lys Arg Ile Ala Gly Ala Leu Leu Glu Arg Gly Leu Pro 165 170 175

Glu Gly Val Asp Met Leu Asn Val Asn Val Pro Ser Asp Ala Thr Glu 180 185 190

Glu Thr Glu Ile Ala Ile Thr Arg Leu Ala Arg Lys Arg Tyr Ser Pro

Thr Val Glu Glu Arg Ile Asp Pro Lys Gly Asn Pro Tyr Tyr Trp Ile 210 225 220

Val Gly Lys Leu Val Gln Asp Phe Glu Pro Gly Thr Asp Ala Tyr Ala 225 230 235 240

Leu Lys Val Glu Arg Lys Val Ser Val Thr Pro Ile Asn Ile Asp Met 245 250 255

Thr Ala Arg Val Asp Phe Glu Glu Leu Val Arg Val Leu Trp Val . 260 270

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 497 AMINO ACIDS
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Met Lys Gly Lys Ser Leu Val Ser Gly Leu Leu Gly Leu Leu Ile 5 10 15

Leu Ser Leu Ile Ser Phe Gln Pro Ser Phe Ala Tyr Ser Pro His Gly 20 25 30

Gly Val Lys Asn Ile Ile Ile Leu Val Gly Asp Gly Met Gly Leu Gly 35 40 45

His Val Glu Ile Thr Lys Leu Val Tyr Gly His Leu Asn Met Glu Asn 50 55 60

Phe Pro Val Thr Gly Phe Glu Leu Thr Asp Ser Leu Ser Gly Glu Val 65 70 75 80

Thr Asp Ser Ala Ala Ala Gly Thr Ala Ile Ser Thr Gly Ala Lys Thr 85 90 95

Tyr Asn Gly Met Ile Ser Val Thr Asn Ile Thr Gly Lys Ile Val Asn 100 105 110

Leu Thr Thr Leu Leu Glu Val Ala Gln Glu Leu Gly Lys Ser Thr Gly 115 120 125

Leu Val Thr Thr Arg Ile Thr His Ala Thr Pro Ala Val Phe Ala 130 135

Ser His Val Pro Asp Arg Asp Met Glu Gly Glu Ile Pro Lys Gln Leu 145 150 155 160

Ile Met His Lys Val Asn Val Leu Leu Gly Gly Gly Arg Glu Lys Phe Asp Glu Lys Asn Leu Glu Leu Ala Lys Lys Gln Gly Tyr Lys Val Val Phe Thr Lys Glu Glu Leu Glu Lys Val Glu Gly Asp Tyr Val Leu Gly Leu Phe Ala Glu Ser His Ile Pro Tyr Val Leu Asp Arg Lys Pro Asp Asp Val Gly Leu Leu Glu Met Ala Lys Lys Ala Ile Ser Ile Leu Glu Lys Asn Pro Ser Gly Phe Phe Leu Met Val Glu Gly Gly Arg Ile Asp His Ala Ala His Gly Asn Asp Val Ala Ser Val Val Ala Glu Thr Lys Glu Phe Asp Asp Val Val Arg Tyr Val Leu Glu Tyr Pro Lys Lys Arg Gly Asp Thr Leu Val Ile Val Leu Ala Asp His Glu Thr Gly Gly Leu Ala Ile Gly Leu Thr Tyr Gly Asn Ala Ile Asp Glu Asp Ala Ile Arg 310 Lys Ile Lys Ala Ser Thr Leu Arg Met Pro Lys Glu Val Lys Ala Gly Ser Ser Val Lys Glu Ser Ser Lys Val Cys Arg Ile Cys Pro Asn Arg Gly Arg Ser Gln Tyr Ile Glu Asn Ala Leu His Ser Thr Asn Lys Tyr Ala Leu Ser Asn Ala Val Ala Asp Val Ile Asn Arg Arg Ile Gly Val Gly Phe Thr Ser Tyr Glu His Thr Gly Val Pro Val Pro Leu Leu Ala Tyr Gly Pro Gly Ala Glu Asn Phe Arg Gly Phe Leu His His Val Asp Thr Ala Arg Leu Val Ala Lys Leu Met Leu Phe Gly Arg Arg Asn Ile Pro Val Thr Ile Ser Ser Val Ser Ser Val Lys Gly Asp Ile Thr Gly Asp Tyr Arg Val Asp Glu Lys Asp Ala Tyr Val Thr Leu Met Met Phe Leu Gly Glu Lys Val Asp Asn Glu Ile Glu Lys Arg Val Asp Ile Asp 475 Asn Asn Gly Met Val Asp Leu Asn Asp Val Met Leu Ile Leu Gln Glu 485

Ala 497

(2) INFORMATION FOR SEQ ID NO:33:

- SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 584 AMINO ACIDS

 - (B) TYPE: AMINO ACID
 (D) TOPOLOGY: LINEAR
- (ii)MOLECULE TYPE: PROTEIN
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Met Pro Arg Asn Ile Ala Ala Val Cys Ala Leu Ala Ala Leu Leu Gly

Ser Ala Trp Ala Ala Lys Val Ala Val Tyr Pro Tyr Asp Gly Ala Ala

Leu Leu Ala Gly Gln Arg Phe Asp Leu Arg Ile Glu Ala Ser Glu Leu

Lys Gly Asn Leu Lys Ala Tyr Arg Ile Thr Leu Asp Gly Gln Pro Leu 50 55 60

Ala Gly Leu Glu Gln Thr Ala Gln Gly Ala Gly Gln Ala Glu Trp Thr
65 70 75 80

Leu Arg Gly Ala Phe Leu Arg Pro Gly Ser His Thr Leu Glu Val Ser

Leu Thr Asp Asp Ala Gly Glu Ser Arg Lys Ser Val Arg Trp Glu Ala

Arg Gln Asn Leu Arg Leu Pro Arg Ala Ala Lys Asn Val Ile Leu Phe

Ile Gly Asp Gly Met Gly Trp Asn Thr Leu Asn Ala Ala Arg Ile Ile

Ala Lys Gly Phe Asn Pro Glu Asn Gly Met Pro Asn Gly Asn Leu Glu

Ile Glu Ser Gly Tyr Gly Gly Met Ala Thr Val Thr Thr Gly Ser Phe

Asp Ser Phe Ile Ala Asp Ser Ala Asn Ser Ala Ser Ser Ile Met Thr 185

Gly Gln Lys Val Gln Val Asn Ala Leu Asn Val Tyr Pro Ser Asn Leu

Lys Asp Thr Leu Ala Tyr Pro Arg Ile Glu Thr Leu Ala Glu Met Leu

Lys Arg Val Arg Gly Ala Ser Ile Gly Val Val Thr Thr Thr Phe Gly

Thr Asp Ala Thr Pro Ala Ser Leu Asn Ala His Thr Arg Arg Gly

Asp Tyr Gln Ala Ile Ala Asp Met Tyr Phe Gly Arg Gly Gly Phe Gly Val Pro Leu Asp Val Met Leu Phe Gly Gly Ser Arg Asp Phe Ile Pro Gln Ser Thr Pro Gly Ser Arg Arg Lys Asp Ser Thr Asp Trp Ile Ala Glu Ser Gln Lys Leu Gly Tyr Thr Phe Val Ser Thr Arg Ser Glu Leu Leu Ala Ala Lys Pro Thr Asp Lys Leu Phe Gly Leu Phe Asn Ile Asp Asn Phe Pro Ser Tyr Leu Asp Arg Ala Val Trp Lys Arg Pro Glu Met 345 Leu Gly Ser Phe Thr Asp Met Pro Tyr Leu Trp Glu Met Thr Gln Lys Ala Val Glu Ala Leu Ser Arg Asn Asp Lys Gly Phe Phe Leu Met Val Glu Gly Gly Met Val Asp Lys Tyr Glu His Pro Leu Asp Trp Pro Arg Ala Leu Trp Asp Val Leu Glu Leu Asp Arg Ala Val Ala Trp Ala Lys Gly Tyr Ala Ala Ser His Pro Asp Thr Leu Val Ile Val Thr Ala Asp 425 His Ala His Ser Ile Ser Val Phe Gly Gly Tyr Asp Tyr Ser Lys Gln Gly Arg Glu Gly Val Gly Val Tyr Glu Ala Ala Lys Phe Pro Thr Tyr Gly Asp Lys Lys Asp Ala Asn Gly Phe Pro Leu Pro Asp Thr Thr Arg Gly Ile Ala Val Gly Phe Gly Ala Thr Pro Asp Tyr Cys Glu Thr Tyr Arg Gly Arg Glu Val Tyr Lys Asp Pro Thr Ile Ser Asp Gly Lys Gly 505 Gly Tyr Val Ala Asn Pro Glu Val Cys Lys Glu Pro Gly Leu Pro Thr 515 Tyr Arg Gln Leu Pro Val Asp Ser Ala Gln Gly Val His Thr Ala Asp Pro Met Pro Leu Phe Ala Phe Gly Val Gly Ser Gln Phe Phe Asn Gly Leu Ile Asp Gln Thr Glu Ile Phe Phe Arg Met Ala Gln Ala Leu Gly Phe Asn Pro His Leu Glu Lys Pro 580

- (2) INFORMATION FOR SEQ ID NO:34:
 - (i) SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 301 AMINO ACIDS
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
 - (ii) MOLECULE TYPE: PROTEIN
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Met Tyr Lys Trp Ile Ile Glu Gly Lys Leu Ala Gln Ala Pro Phe Pro 10 15

Ser Leu Gly Glu Leu Ala Asp Leu Lys Arg Leu Phe Asp Ala Ile Ile 20 25 30

Val Leu Thr Met Pro His Glu Gln Pro Leu Asn Glu Lys Tyr Ile Glu
35 40 45

Ile Leu Glu Ser His Gly Phe Gln Val Leu His Val Pro Thr Leu Asp 50 55

Phe His Pro Leu Glu Leu Phe Asp Leu Leu Lys Thr Ser Ile Phe Ile 65 70 75 80

Asp Glu Asn Leu Glu Arg Ser His Arg Val Leu Val His Cys Met Gly 85 90 95

Gly Ile Gly Arg Ser Gly Leu Val Thr Ala Ala Tyr Leu Ile Phe Lys
100 105 110

Gly Tyr Asp Ile Tyr Asp Ala Val Lys His Val Arg Thr Val Val Pro 115 120 125

Gly Ala Ile Glu Asn Arg Gly Gln Ala Leu Met Leu Glu Asn Tyr Tyr 130 135 140

Thr Leu Val Lys Ser Phe Asn Arg Glu Leu Leu Arg Asp Tyr Gly Lys 145 150 155 160

Lys Ile Phe Thr Leu Gly Asp Pro Lys Ala Val Leu His Ala Ser Lys

Thr Thr Gln Phe Thr Ile Glu Leu Leu Ser Asn Leu His Val Asn Glu
180 185 190

Ala Phe Ser Ile Ser Ala Met Ala Gln Ser Leu Leu His Phe His Asp 195 200 205

Val Lys Val Arg Ser Lys Leu Lys Glu Val Phe Glu Asn Met Glu Phe 210 215 220

Ser Ser Ala Ser Glu Glu Val Leu Ser Phe Ile His Leu Leu Asp Phe 225 230 235

Tyr Gln Asp Gly Arg Val Val Leu Thr Ile Tyr Asp Tyr Leu Pro Asp 245 250 255

Arg Val Asp Leu Ile Leu Leu Cys Lys Trp Gly Cys Asp Lys Ile Val 260 265 270

Glu Val Ser Ser Ser Ala Lys Lys Thr Val Glu Lys Leu Val Gly Arg

> 280 285

Lys Val Ser Leu Ser Trp Ala Asn Tyr Leu Asp Tyr Val

- (2) INFORMATION FOR SEQ ID NO:35:
- (i) SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 257 AMINO ACIDS
 (B) TYPE: AMINO ACID

 - (D) TOPOLOGY: LINEAR
- MOLECULE TYPE: PROTEIN (ii)
- SEQUENCE DESCRIPTION: SEQ ID NO:35: (xi)

Met Arg Ile Leu Leu Thr Asn Asp Asp Gly Ile Tyr Ser Asn Gly Leu

Arg Ala Ala Val Lys Gly Leu Ser Glu Leu Gly Glu Val Tyr Val Val

Ala Pro Leu Phe Gln Arg Ser Ala Ser Gly Arg Ala Met Thr Leu His

Arg Pro Ile Arg Ala Lys Arg Val Asp Val Pro Gly Ala Lys Ile Ala

Tyr Gly Ile Asp Gly Thr Pro Thr Asp Cys Val Ile Phe Ala Ile Ala

Arg Phe Gly Asp Phe Asp Leu Ala Val Ser Gly Ile Asn Leu Gly Glu

Asn Leu Ser Thr Glu Ile Thr Val Ser Gly Thr Ala Ser Ala Ala Ile

Glu Ala Ser Thr His Gly Ile Pro Ser Val Ala Ile Ser Leu Glu Val

Glu Trp Lys Lys Thr Leu Gly Glu Gly Glu Gly Ile Asp Phe Ser Val

Ser Ala His Phe Leu Arg Arg Ile Ala Thr Ala Val Leu Lys Lys Gly

Leu Pro Glu Gly Val Asp Met Leu Asn Val Asn Val Pro Ser Asp Ala

Ser Glu Gly Thr Glu Ile Ala Ile Thr Arg Leu Ala Arg Lys Arg Tyr

Ser Pro Thr Ile Glu Glu Arg Ile Asp Pro Lys Gly Asn Pro Tyr Tyr

Trp Ile Val Gly Arg Leu Val Gln Glu Phe Glu Pro Gly Thr Asp Ala

Tyr Ala Leu Lys Val Glu Arg Lys Val Ser Val Thr Pro Ile Asn Ile

225 230 235

Asp Met Thr Ala Arg Val Asp Phe Glu Asn Leu Gln Arg Leu Leu Ser 250

Leu

- (2) INFORMATION FOR SEQ ID NO:36:
 - $(\dot{\mathbf{1}})$ SEQUENCE CHARACTERISTICS
 - (A) LENGTH: 264 AMINO ACIDS (B) TYPE: AMINO ACID

 - (D) TOPOLOGY: LINEAR
 - (ii) MOLECULE TYPE: PROTEIN
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Met Glu Asn Leu Lys Lys Tyr Leu Glu Val Ala Lys Ile Ala Ala Leu

Ala Gly Gly Gln Val Leu Lys Glu Asn Phe Gly Lys Val Lys Lys Glu 20 25 30

Asn Ile Glu Glu Lys Gly Glu Lys Asp Phe Val Ser Tyr Val Asp Lys

Thr Ser Glu Glu Arg Ile Lys Glu Val Ile Leu Lys Phe Phe Pro Asp

His Glu Val Val Gly Glu Glu Met Gly Ala Glu Gly Ser Gly Ser Glu

Tyr Arg Trp Phe Ile Asp Pro Leu Asp Gly Thr Lys Asn Tyr Ile Asn 85 90 95

Gly Phe Pro Ile Phe Ala Val Ser Val Gly Leu Val Lys Gly Glu Glu

Pro Ile Val Gly Ala Val Tyr Leu Pro Tyr Phe Asp Lys Leu Tyr Trp

Gly Ala Lys Gly Leu Gly Ala Tyr Val Asn Gly Lys Arg Ile Lys Val 135

Lys Asp Asn Glu Ser Leu Lys His Ala Gly Val Val Tyr Gly Phe Pro

Ser Arg Ser Arg Arg Asp Ile Ser Ile Tyr Leu Asn Ile Phe Lys Asp

Val Phe Tyr Glu Val Gly Ser Met Arg Arg Pro Gly Ala Ala Ala Val

Asp Leu Cys Met Val Ala Glu Gly Ile Phe Asp Gly Met Met Glu Phe

Glu Met Lys Pro Trp Asp Ile Thr Ala Gly Leu Val Ile Leu Lys Glu

Ala Gly Gly Val Tyr Thr Leu Val Gly Glu Pro Phe Gly Val Ser Asp

225 230 235 240

Ile Ile Ala Gly Asn Lys Ala Leu His Asp Phe Ile Leu Gln Val Ala 245 250 255

Lys Lys Tyr Met Glu Val Ala Val 260

Pyrolobus fumarius 1A (1ph7) SEQ ID NO.37

1 TGC CCG AGC GTG TTG CCA AGA TGC TTG AAA GAA TGC TAT CCA AGG CGG AAT CTA TGC TCG 61 GCG ACG CCC AGA GGC TTA TCG AGG AGG GTA AGG CCG TTG AGG CTA AGA AGC TGT TAG CGG 121 CTG CTC ATA GGC TAG TAG ATC GCC TAG AGG ATG CTC TCG ACC ACG CCC TCA ACC ATA TAG 181 AGC ATC AGA AGG AAC ATC ATG AGG AGC ACC ACA AGG AGC ACG ACT AAC AAC ACT CTT AGA 241 ATC TCG AGA CGA GCT TGC TTC CCG TGT CTC TCG CGC CTA GCC AGT TTT TAA TAG CCT AAG 301 CCG AGA CCC ACA TTC CAA CAT TAC TCC GTT TGT CAC TAT CAT GTT CTA ATT GTC ACA CGC 361 CCC GTA TAA ATT GGG GGA CCT GGA GGA AGC GTT GCC GGT GAC CCC GCG TGG CCA AGA AGG 421 CTG TCT GCC CAA TAT GCG GTG GCG ATG TTG AAC TAC CCG ATA ACG TAA TGG ATG GCG AGA 481 TCG TGG AGC ACG ACT GTG GGG CAA TGC TAG TCG TGA GGA TCC GGG ATG GCA ATG TTG TTC 541 TAG AGC AGT TGG AGC GCG TTG AGG AGG ACT GGG GAG AGT AGA GGC TAT GCG CAT AGC AAT 600 601 CGT TTA TGA CCA TCC GCG TGT TGA GGA GAA GAG GTT AGC TGA GGA AGC GAA GCT TGG 661 TCA CGA ACC TGT CCT CTT TAA TAT TGA CTC GTT GCT CTT TCG CCT TGA TAG CCT GGA GCG 721 CAT TCT AGG CGA TGT TGA TGT AGT ACT TCA GAG GGC GGT GAG TTA CTT CAA GGC TCT CGA 780 781 GTC TAC AAG GAT ACT CGA GGC TGC CGG CTA CAC TGT CAT CAA CAA TAG TTT AGT GCA GCT 841 TAA CTG CGG CGA CAA ACT ATT GAC AAC GAT CTT GCT TGC TAA GCA TGG TGT GCC AAC ACC $90\overline{1}$ GCG TGC ATA CGC TGC TTT TTC GCG TGA CAC TGC TGT GCG GGC TGC AGA GGA GCT TGG ATA 960961 CCC CGT TGT TGT CAA GCC CGT CAT TGG TAG TTG GGG TAG GCT TGT GGC TAG GGC TGA TTC 1020 1021 CAG GGA GAG TCT AGA GGC TGT GAT AGA GCA TAG AGA GGT TCT CGG CCC GGC TTA CTA CAA 1081 GGT TCA TTA TGT GCA AGA GTA TGT GCG CAA GCC TCT ACG TGA CAT ACG CGT ATT CGT GAT

1141 1200	TGG	TGA	TGA	ggt	TCC	CGT	GGC	GAT	ATA	CAG	GGT	TAA	CGA	GCG	TCA	TTG	GAA	GAC	TAA	CAC
1201 1260	GGC	ACT	AGG	CGC	CAA	GGC	CGA	GCC	TGC	GCC	AGT	GAC	ccc	CGA	GTT	ACG	TGA	GTT	AGC	GCT
1261 1320	TCG	CGC	GGC	CAA	GGC	ŢĠŢ	GGG	TGG	CCG	TGT	GCT	TGG	TAT	AGA	TGT	GTT	TGA	AGA	ccc	GGA
1321 1380	GAG	AGG	CCT	CCT	CGT	GAA	CGA	GAT	TAA	CGC	gaa	CTC	GGA	CIT	CAA	GAA	CAC	TGA	GAG	GGT
1381 1440	GAC	CGG	GTT	TAA	CAT	GGC	TAG	GGC	TAT	CGT	CGA	GTA	TGC	AGT	GTC	GGT	cgc	GAA	GAG	GTG
1441 1500	AAT	GGA	ATG	GAT	AGG	GTA	GAG	GTG	CTT	CTG	GAT	GAG	GCT	AGG	CGT	GGC	GCT	ATA	GAG	GGT
1501 1560	GAC	GCT	CGC	CGC	GCA	TGT	GAA	GCG	GCA	TTA	AGG	CTG	GTT	GAC	GTT	GTG	CTC	CGC	GAG	GGG
1561 1620	CCT	AGG	GTT	GCA	CAG	GAG	TCT	GGG	CGT	GGG	ATT	gaa	ccc	ggt	GAT	gta	CTA	CTA	GCT	GAG
1621 1680	GCT	CTG	AGC	TTG	AGA	GCA	GAG	CAG	GTG	AAG	GAG	GAG	ccc	AAG	GCG	GAC	AAT	TGT	CTG	GAG
1681 1740	CTC	GCA	AAG	GCT	GCA	TTC	CGC	CTC	TAT	AAG	CGG	CTC	CAG	GGG	ATG	GAG	TAA	AGT	TCG	CAG
1741 1800	TGT	GTT	GCC	CGT	TTT	AGC	crc	TGC	CIT	ACT	TTC	TAC	TCG	CGT	GAG	GCG	AGT	GTC	CCT	TGA
1801 1860	CAC	GTT	GCT	GGC	GCG	AGC	TGA	GAA	ACG	ACC	TCG	AGA	TGA	TAC	ccg	AGA	TCG	TCG	aga	AGC
1861 1920	AGA	TCG	AGG	AGA	CGA	TAG	TGC	CGG	AGG	GTC	TTG	GCG	AGC	AAC	GAC	TTG	TGT	TCA	TTG	GCA
1921 1980	GCG	GTG	ATT	CTT	TCG	CGG	CCG	CAC	TTG	TAG	CCG	AGC	atg	CCG	gca	TAG	GCG	TCG	CAC	GCG
1981 2040	ATC	CTC	TTG	ATG	TGC	TAG	TGG	CTG	GCG	TTG	ATG	GGC		GCG	ACG	CTA	TAC	TCC	TAA	GCG
2041 2100	TTG	GT G	GGC	GCT	CAA	AAC	GAG	TTG	TTG	ACG	CGG	CTC	 GTT	TCC	TGT	CTT	CAC	GTG	GCT	TTC
2101 2160	GTA	TCA	TAG	CGG	TCA	CGG	GTA	ACG	aga	GGA	GTC	CTC	TCG	CAC	GCA	CAG	CAC	A CG	TTA	CCG
2161 2220	TGA	AGC	TCG	TCT	ATT	CTG	ACC	TCG	CCT	GTG	GCA	TGG	GCG	ccg	CAC	GCC	ATG	TCG	CTA	TGC
2221 2280	TTG	CAG	CGC	TCT	ccg	CAT	TGT	TCA	ACG	CTA	GAC	ctc	GTA	TAC	ccg	AGA	AGC	TTG	TTG	AGG
2281 2340	AGC	ccc	TGC	CTT	TCG	ACC	CTC	AGG	ctg	TGT	ACG	CGG	GTG	TGG	GCG	TTG	GTG	TAG	CCT	CTG

2341 2400	ccc TGT	TCA	TGG	TGT	TGA	AGA	TCT	GCG	agt	TGC	TCG	CAG	ACT	GCG	CCA	CCT	GGT	GGC	ATC	-
2401 2460	TAG AGC	AGT	TCG	CAC	ACG	CAC	CTG	TCT	ATG	GCA	CGA	gaa	GCA	ATA	TAC	TCG	TCG	TGT	ATC	
2461 2520	CGA TCC	TCG	TTG	TGA	GAG	GAG	CAC	GCT	AGA	GGA	GTA	TCT	CTC	GGC	CTT	CCG	GGA	GGC	CGG	
2521 2580	GTT TGA	GGT	CAC	CAC	TGT	ACC	CGT	GTT	GAA	CGA	ccc	TTG	GTC	TAC	AGC	TAT	TCT	CCA	CGC	
2581 2640	TAC GCT	GGC	CAT	CTC	CAG	TGC	TGC	AGA	GAC	CGC	CTT	CAG	TCG	CGG	CAT	TGA	GGA	GCC	GGG	
2641 2700	ATA TCG	TGC	ACA	TCC	CGC	GCT	TAG	CAG	GCT	AAC	CAG	GCT	GAT	CTA	CCT	AGA	GGA	GTA	GAA	
2701 2760	cci cic	GAG	GAC	CGG	TAT	GTA	GTG	GTC	TAG	AGG	CTT	ccc	GTC	ATG	GTG	TAT	CGC	GAG	GCC	
2761 2820	TAT TCC	TGC	TCT	CCT	CGC	GCC	TTC	CAC	gtt	GGG	CTC	ATA	ATC	ATC	TAT	GAA	TGC	TGT	TTT	
2821 2880	CGC TGG	GTC	CGC	GCG	AAG	GAG	TTG	CAT	CGC	CGC	CTC	gta	TAT	CTT	TGT	GTG -	T G G	CTT	GCA	
2881 2940	AAA GCC	GAC	AAT	ATC	CCT	CGT	AAC	CAC	CGT	ATC	CAC	GAG	GTG	GGC	TAG	ATC	GTC	ACG	CTC	
2941 3000	TAG AAG	TAG	ACG	TAC	GCA	TTC	GTA	GCA	CCA	GTT	GTT	CGA	GAC	TAT	GCC	GAC	CAG	TAT	ccc	
3001 3060	GTT TCT	CTT	GGC	CCA	TCT	TAG	CAG	CTC	GTA	TGT	ACC	CGG	TGC	TAC	GTA	TAC	GCC	AGA	CAG	
3061 3120	CAC AGC	TGA	TTG	CAA	TAC	CCT	TGC	TAA	TGC	CTC	TGC	CCT	TGA	GGG	GGT	CGG	CGT	CAA	GCC	
3121 3180	GTG TTT	TGC	GAG	GAG	CAC	GGC	AGC	CCC	ATA	CAC	TAT	ACT	TTG	TTG	CAC	GGA	GAC	ATC	CAG	
3181 3240	CCT CCA	CGT	GTC	CAT	TAC	ACG	CCT	CAC	GCT	ATC		CGT	CGC	GTC	GGC	ccc	TAG	GGC	ACG	
3241 3300	TAG ATG	TCT	GGC	AGC	AGT	CTC	GTA	GAG	AGT	CTC	CIC	GTA	CCA	CTC	ATT	TGT	GAG	GTA	AAT	
3301 3360	GAC GCC	ACC	TAA	ATC	CAG	CAG	GAG	TGT	AGG	GTT	ACG	CGG	CAA	GGC	GCC	TCC	TCA	TGT	ATT	
3361 3420	CGA GGA	GGC	CGC	CCG	TTG	CCA	GAA	TTT	CAG	CTA	CAA	CAC	ccc	GGA	AGG	GCG	GGA	AAC	GGT	
3421 3480	ACG TCA	. ACA	CCC	TAC	CAT	CCT	TCT	TGA	TGA	GCT	TCG	CTA	CAC	CCT	CGT	CAA	GGT	TTA	TCT	
3481 3540	CTA TCT	CGT	CGC	CCT	CCT	CGG	CCG	CCT	CCA	CGA	GCT	CTG	GGA	GCA	CTA	TAA	CGG	GGA	GCC	

3541	CGT TGT	TAA	TCG	CGT	TAC	GGT	AGA	ATA	TTC	TCG	AGA	AGC	TCT	TCG	CTA	TGA	TGG	CCT	TGA	
3600																				

- 3601 CGC CTG CAG CCT TGA GAG CTA TCG CGG CTT GCT CCC TGC TAC TAC CCA TAC CAA AGT TCC
- 3661 TAC CCG CGA CCA GCA CTA CAC CCT TGG ACG CCT TCT TGG GGA ACT CCG GAT CCA GAG GCT
- 3721 CCA TAG CAT GCT CGG CAA GCT TCT CCG GCT CAG TAT ATA CCA GGT AGC GGG CAG GGA TAA 3780
- 3781 TCA CGT CGG TGT TGA TGT TAT TGC CGT AAT TGA GCA CAG GGC CCT TCA CGA CAC CCA GGT 3840
- 3841 TCA AGA GAG GTT CAC CAC AAG TTT GGC CTC GCT ATC CCA GGC TAT AAT CCA GCT GTT TAC
- 3901 TCG GCC AGC TTC ACC CAC ACA CTT TTC AAC TCC ATT ATC CTT GTA GCG CAA TCT ACC CTT
- 3961 CTG GGT AGC ACA GCG TTA AGC CCA TAG TGC CAA GGC GCC ACA ATG ATG CCC TCC GGC ACA
- 4021 TTC TCG TCG GGT ATC AGC CGG AGG CGT ATG GCC CCT CTC TCC GTC TCG AGC CTA GCG TGA
- 4081 CCG GCG CCA GCC TCC TTA GGG TTG ACT CGT GCG TAT AGC TCG CCG CTC ACA TCT AGC ATC 4140
- 4141 GCG TTT GTA CAG TAG CTC ACC GGG TCT CTT GCA GTC ACG AGC ACC TTC CTA TCA CCA TCG
- 4201 GGC ACG ACC GGC TCG ACC GGC GGG TAT AGA CGG ACG CGT ATC CTC GAG ACA CGC CTG GGC 4260
 - 4261 AGG AGG TAC TCG CCT CTC TCC GCA ACC GCC TTG GAG GAA 4299

-76-

Thermococcus 9N-2 (31ph1)

SEO ID NO:38

1 TGG ACT GAT AAA GAA AAA GAA GAG GTT TAA GGG CCT CAA TAT TAA ATT CTA CAC ATT AGA 61 TAT CCA AAA TGG AGA ATT ACT TAA TCT AGA GAC TTA CCT TAA GGA GTT ACA TGA GTT CCT 121 TAG AGG CCT TAC ATT AAA ACG AAA AGT AGA AGA GGA ACA ATG ACC CCC GAA GAG CTC CTA 181 ACC CGC CTC GAA TTC AAA GGA GTA ACC CTC GAA AAG ATG CTC AAT ACT GCG TTA GAG CTC 241 TAC ATC GGC GAC GAG CGC GAG AAA GTT CGA GAA AGG CTG AGA GAG CTG ATG CTG AGG TAT 300 301 CTG GGC GAC ATC AAC GTT CAA GCT CTG CTC TTT TCG GCT CTA CTG CTC GAA GAG AAC TTC 361 AAG GTT GAG GGC GAC CCC GTG AAC CTT GTG GCC GAC GAG CTC ATC GGC ATG AAC ATC GCC 421 GAG CTC ATA GGT GGA AAG ATG GCG CTC TTC AAC TTC TAC TAC GAC ACC AAG AAG CCC 480 481 GGC ATT TTA GCC GAG CTT CCG CCT TTC CTC GAC GAT GCG ATA GGG GGC TTT ATA GCG GGC 541 TGT ATG ACA AGG CTG TTC GAG GGG GTG TAC GGT GCG GAA TCT CTT ACC CTT CTT CAC GCG 601 GAT TCC GGT CAA AGG CAA CTT CAA AAG GGT TAG AAA TGA GCT CTG GGC ACT TCC CAT TCT 661 CGC ACC GGT AAC TTC GGC CCT GGC GAC GCT CGT GGG CTC TGT GCT CGC CGG GGT AAT AAT 721 CCT GGG CGG CAA CTA CGC GTT TCA CCC AAC GTC TCG GCA ACC CAC GTG CTG ATA ACC CTC 781 ATA GGC TTC GTC GTG GTC TAC AGC ATA CTG TTC TAC ATC TGG CTC CAC TTC GTC AGG AAG 841 CTC ATC AGG GAG GGC CCC GAA CCG GTT GAG GGT GAC GTC ACC GCG AAG CCG ACC CCT GCC 901 GTT AGC GCC GCG GGA GGT GGT CAG TGA TGG ACT ACG CGA CCG CAT GGT TTT ACT TCT CCG 961 CCT TCC TCC TCG GAA TGT ACT TAG CGT TTG ATG GCT TCG ACC TTG GCA TAG GCG CGT TGC 1021 TCG CCC TGA TTA AGG ACC AGA GGG AGC GCG ACA TAC TCG TGA ACA CCA TCG CGC CGG TCT 1081 GGG ACG GCA ACG AGG TCT GGT TCA TCA CCT GGG GTG CCG GGC TCT TCG CGA TGT GGC CGG

1141 1200	CGC T	CT ;	ACG	CGA	CGC	τςτ	TCA	GCA	CGT	TCT	ACC	TTG	CCG	TCT	GGC	TGC	TCG	CGT	TCC	TGT
1201 1260	TCA T	AT 3	rca	GGG	CTG	TCG	GCT	TTG	AGT	TCA	GGA	ACA	AGA	ACA	AGG	AGC	TAT	GGG	ACA	AGC
1261 1320	TCT T	23 (Ξτc	TCG	TCA	GCG	CGT	TAA	TCC	CGC	TCG	TCA	тсс	GCG	TCA	TAG	TCG	GCA	ACC	TCA
1321 1380	TCA TO	GG (AAG	TTC	CCA	TTG	ACG	CCA	AGG	GCT	TCC	ACG	GCT	CAC	TGC	TGA	cgc	TCT	TCA	GGC
1381 1440	CCT A	CC (CGC	TCA	TCG	TCG	GCC	TCT	TCA	TAC	TCT	TCG	CGG	TGA	CCT	GGC	ACG	GAG	CCA	ACT
1441 1500	GGG G	CG 7	CT	ACA	AAA	CCA	CAG	GAA	AGC	TCC	AGG	AGC	AGA	TGA	GGG	AGC	TCG	CCT	TCA	AGG
1501 1560	CCT G	SC 3	cc	TGA	CCG	TCG	TCT	TCC	TCC	TGC	TCA	CAG	TCA	TCG	GCA	TGA	AAA	TCT	GGG	ccc
1561 1620	CAC TO	GA (GT	TCG	AGA	GGG	CAC	TAA	CGC	CGC	TTG	GGC	TCC	TCC	TAA	CGG	TTG	TCA	TCC	TCG
1621 1680	TGG C	AG (SAC	TGC	TCG	ACG	GAC	AGC	TCA	TCA	AGA	AAG	GGG	AGG	AGA	ATT	TGG	CCT	TCT	ACA
1681 17 4 0	TCA G	CT C	SGC	TGG	CCT	TCC	CGC	TCG	TTG	TGT	TCC	TCG	TCT	ACT	ACA	CAA	TGT	ACC	CCT	ACT
1741 1800	GGG T	CA 7	rct	CGA	CCA	CCG	ATC	CGA	ACT	TCA	AGC	TCA	GCA	TAC	ACG	ACC	TCG	CGG	CAT	CTC
1801 1860	CGC T	GA (ccc	TCA	AGG	CCG	TCT	TGG	GAA	TCT	CGC	TGA	TCC	TGG	CGG	TCA	TCA	TCA	TGG	cci
1861 1920	AÇA C	ככ ז	rcr	ACG	TAT	ACA	GGG	CCT	TCG	GCG	gaa	AGG	TCA	CCG	AGG	CGG	AGG	GCT	ACT	ACT
1921 1980	GAG T	rc	ccc	TTT	CCT	TTT	TCG	ATA	TTC	GAA	CTT	TTT	TAG	GGA	AAA	GTT	TAT	AAT	TCG	AGT
1981 2040	CAC C	TA A	AGT	TCC	TTC	TGG	AAA	CCT	AAA	AAA	CGG		TCG	AAA	TGC	ACA	GAG	GCA	GAT	CTA
2041 2100	CCG G	CT (GGC	CCT	ACG	ACC	GGA	AGC	CGG	TCC	TCG			GGG	AAA	CCA	CCA	AAG	CCI	GCC
2101 2160	GGC T	CA A	agt	GCA	AGC	act	gca	gag	CGG	AGG	CAA	TAC	TCC	AGG	CAC	TGC	CGG	GCG	AGC	TGA
2161 2220	ACA C	GG J	AGG	AGG	GAA	AGG	ccc	TCA	TCG	ATT	ccc	TCA	CCG	ACT	TCG	GAA	GGC	CCT	ACC	CGA
2221 2280	TAC T	CA T	rrc	TCA	ccg	GTG	GCG	ACC	CGC	TCA	TGA	GGA	AGG	ACA	TCT	TCG	AGC	TCA	TCG	AGT
2281 2340	ACG C	CG :	TTG	AGA	AGG	GCA	TTC	GCG	TTG	GTC	TCG	ccc	CCG	CTG	TAA	CGC	ccc	TCC	TGA	ccg

2341 2400	AGG AA	A CAA	TCG	AGA	GAA	TCG	CGA	GGA	GCG	GAG	TTA	agg	cgg	TAA	GCA	TAA	GCC	TCG	ACA
2401 2460	GCC CG	T TTC	CAG	AAG	TTC	ACG	ACG	CAA	TCA	GAG	GCA	TAG	AAG	GGA	CGT	GGG	AGA	AAA	cca
2461 2520	TCT GG	G CCA	TĆA	AGG	AGT	TCC	TGA	AAC	ACG	GCC	TAA	GCG	TTC	AGG	TGA	ACA	CGG	TTG	TGA
2521 2580	TGC GC	g aga	CCS	TTG	AAG	GAC	TGC	CCG	AGA	TGG	TGA	AAC	TGC	TTA	AAG	ACC	TCG	GCG	TCG
2581 2640	AAA TO	T GGG	AGG	тст	TCT	ACC	TCG	TCC	CGA	CCG	GGA	GGG	GCA	ACT	TCG	AGA	GCG	ACC	TGA
2641 2700	GGC CG	G AGG	AGT	GGG	AGG	ACG	TCA	CAC	ACT	TCC	TCT	ACG	AGG	CCT	CGA	AGC	ACC	TCC	TCG
2701 2760	TGA GG	A CCA	ccg	AGG	GCC	CGA	TGT	TCA	GGC	GAG	TGG	CGA	TAA	TGA	GGA	AAG	ccc	TTG	AGG
2761 2820	AGA AG	g gat	TCG	ACC	CCG	ACG	AGG	TTC	TCA	AGC	CCG	GGG	AGC	TCT	ACT	TCC	GGC	TGA	AGA
2821 2880	AAC GO	C TCG	TTG	AGC	TTC	TCG	GCG	AGG	GGA	ACG	AGG	CGA	GGG	CCC	AAA	CTA	TGG	gaa	CGC
2881 2940	GCG AC	g gga	AGG	GAA	TAG	TCT	TCA	TCG	CCT	ACA	ACG	GCA	ACG	TCT	ACC	CGA	GCG	GTT	TCC
2941 3000	TGC CC	T TCA	GCG	TCG	GCA	ACG	TCC	GCG	AGA	ааа	GTT	TGG	TTG	AGA	TTT	ACA	GGG	AGA	GTG
3001 3060	AAC TT	A TGA	AAA	AGC	TCC	GCT	CGG	CCG	AGT	TCG	AGG	GGC	GCT	GCG	GGA	GGT	GCG	AGT	TCA
3061 3120	GGG A	A TCT	GCG	GGG	GAA	GCA	GGG	CGA	GGG	сст	ACG	CCT	ATC	GCT	TAA	ACC	cgc	TÇG	CCG
3121 3180	AAG AG	c crg	CCT	GCC	CGT	ACG	AGC	CGG	GCT	CAT	ACC	TAA	GGC	TCG	CCA	ааа	AGT	TCA	ATC
3181 3240	TTC AC	C TTC	CGA	TTG	AGA	TTT	TTG	GAG	ccc	AAA	AGC	CGA	TTT	GAG	GTG	ATG	GAA	ATG	AGG
3241 3300	TGG AA	G GCT	GTT	TTA	CTG	ATT	GGA	ATC	CIC	CIC	GTG	TCT	GTC	CTC	GGT	GCC	GGA	TGC	GTT
3301 3360	GGC TO	G AAT	ACC	TCA	ACT	GAA	ACC	GGC	CCA	TCC	CAG	AAG	GAA	ATA	ACC	GTG	AAG	GAC	TTC
3420	TCG GC	A AGG	AAC	ATC	ACG	GCT	AAA	GTT	CCG	GTT	CAG	CGG	GCG	GTC	GTT	CTC	TCG	ACT	TCC
	TCG GC																		

3541 3600	CTC	AAG	ATT	GAC	GAC	TGG	GAG	AAG	GTT	TTA	GCC	CTA	AAG	CCC	GAC	CTA	ATC	ATC	GAC	CTC
3601 3660	GAC	CTG	AAG	AAG	TTC	TAC	AAC	GTT	GAC	GAG	CTC	CTC	AAC	CGC	TCC	GCC	AGC	TAC	GGA	ATT
3661 3720	CCG	GTC	GTC	CTG	CTG	AGG	GAG	GAT	AAC	CTT	GAG	GAC	ATA	CCG	AAG	GCG	GTT	TCG	CTC	CTC
3721 3780	GGT	CAG	CTC	TTC	GGA	AGG	GAG	AAA	GAG	GCC	AAG	GCC	TTC	GAC	GAC	TAC	TTC	AAC	GAG	CAG
3781 3840	GTG	AAG	GAG	GTT	AAG	GCC	ATA	GCC	TCA	AAG	ATT	CCA	GCG	GAG	GAG	AGA	AAG	AAG	GCG	ATA
3841 3900	ATG	ATA	CAG	CCG	ATA	ATG	GGC	AAG	стс	TAC	CTC	GTC	AAC	GGC	AAC	GAC	GTC	CTT	GCT	CAG
3901 3960	GCC	GTC	AGG	CTC	GTT	GGG	GCG	GAC	TAC	CTC	GTG	AAC	CTG	ACC	TTC	Aac	GGC	TAC	act	CCG
3961 4020	GTT	AGG	GTC	CCG	ATG	GAC	GGG	GAG	AAG	ATA	ATA	GCG	AAC	TAC	CGC	GAT	GCA	GAC	GTC	GTA
4021 4080	ATC	CTC	CTG	ACG	AGC	GCC	GTA	ACG	CCT	TAC	GAC	CAG	GTC	GAG	AAG	CTC	cgg	GAG	GAG	ATG
4081 4140	CTC	AGC	GAC	GAG	GCC	TGG	AGG	GGC	ATT	AAG	GCC	GTC	AGG	GAG	GGC	AAC	GTA	GTA	ATC	CTC
4141 4200	AGG	GCG	GAC	ATG	GGT	AAA	GAC	TCC	TTC	CTC	CGC	TGG	AGC	CCG	ccc	TTG	GCA	GTG	GGA	ATO
4201 4260	TGG	GTC	ATT	GGA	AAG	GCA	ATC	TAC	CCG	GAC	TAC	TAT	CCT	GAC	TGG	AAC	GAC	AAG	GCC	AAG
4261 4320	GAC	TTT	CTG	AAG	AGG	TTT	TAC	GGC	CTC	TCC	TGA	TTT	TTC	TTT	TGG	GGT	GGG	ACG	ATG	ATA
4321 4380	GCG	GTC	TTT	CCA	GCG	AGT	CTC	GCG	GAA	ATC	GTC	AAA	CTC	GTC	ÇGG	AAA	GCC	GGG	GAG	ATA
4381 4440	GCC	GGA	GTG	AAC	GAG	GAA	ATC	AGG	TTC	GAC	ccc	TGC	CTG	CCG	GAG	CTG	AAG	GAT	AAG	CCT
4441 4500	GTC	ATC	GGA	AAG	TAC	CTC	AAG	CGG	AGC	AAG	AGG	ACC	TAC	TGG	GAC	GTT	TTA	GAG	GAG	CTI
4501 4560	AGG	CCG	GAC	CTT	ATC	CTC	GAC	TTC	GAT	GTT	GAG	AAC	CIG	CAC	TCC	GGG	GAC	GAG	CTG	AGG
4561 4620	GCC	TTT	GGG	GAG	CGT	ATA	GGG	GCA	AGG	GTC	GAG	CTG	ATT	GAC	TTC	GAG	ACC	GTT	GAA	GGC
4621 4680	TTC	GTC	gag	GCG	AGC	AGG	AGG	ATA	GCC	GAG	CTA	ACG	AGG	GGC	GAC	TTT	TCA	AAG	CTC	GGC
4681 4740	GGG	TTC	TAT	GAG	AAG	CAC	CTG	ACG	AGG	CTG	GGT	GAG	ATA	ACT	GAA	GCC	ATC	GAG	GAG	AGC

4741 4800	CCT AAA	GCC	CIG	CTC	ACC	TAC	CGG	A AC	TTC	AAC	GTC	GTA	ACG	AGG	ACC	AAC	GT T	CTG	AGC	
4801 4860	GAC GCG	GTT	AGA	AAA	GCA	GGG	GCG	ATG	AAC	CTC	GGC	GAG	AGG	ATA	CGG	ACA	AAG	CGG	AAG	
4861 4920	GTC TAT	ccs	GTA	AAG	AAG	GAG	cgc	TTC	TTC	AGG	TCC	TTC	GGC	GAT	GCG	GAG	CAC	CTC	TTC	
4921 4980	CTG CTC	ACG	AGC	ATA	ATG	ACG	GAC	AGG	GAG	AAA	ATG	GAG	GGG	ATA	AGG	GAT	GAA	ATC	CIT	
4981 5040	GAC TCG	GCC	GAG	TGG	AGG	GCA	ATG	GAA	GCC	GTT	CAG	CTC	GGA	AAC	GTG	CAC	ATA	GTT	GGC	
5041 5100	TCG GCC	CTC	GAC	CTT	GAG	AGC	TTC	ATG	CGC	TGG	agt	ccc	cgc	ATA	ATC	CCG	GGA	ATC	TAC	
5101 5160	CAG CTT	GGA	AGG	TTT	ATA	CAC	GGA	ACA	AAT	CAC	CCA	CGA	ATC	TCG	TGG	AAA	TCA	CTG	CAA	
5161 5220	AAG TTT	AAA	ATC	ccc	стс	CCA	CCC	CTC	GAA	GAA	CAA	AAA	CGC	ATC	GTC	GCC	TAC	CTC	GAC	
5221 5280	TCG ATA	CAC	GAG	CGC	GCC	CAA	AAG	CTG	GTA	AAG	CTC	TAC	GAG	GAG	CGG	GAG	AAG	GAG	CTT	
5281 5340	GAG AAG	CTT	TTC	ccc	GCG	GTG	CTT	GAT	AGG	GCG	ттт	AGG	GGT	GAG	CTG	TGA	TTC	CGG	GAA	
\$341 5400	TGG AAT	ACG	GCT	TTG	AGA	GGG	CAA	TCT	TTG	AGA	TAG	TCA	GCG	GCT	TTG	TTC	TCT	ccc	TCG	
5401 5460	TAG TCA	GGG	CTT	TCG	CTT	ACA	GTT	TTG	GTC	TTC	CAT	GGG	TAT	CCT	TTT	TGT	TCA	ACG	TTC	
5461 5520	TTT CGA	TAC	TTC	TGA	CAA	TAG	GCC	TGA	TTG	ACA	AAA	TGC	CCT	TCT	GGT	CCA	TGT	CAT	ATC	

-81-

OC1/4V (33ph1)

SEQ ID NO:39

1 AGC TTG GAT ATC GAA TTC CTT ATA TGA AAA ATT CAT CGA ATT GGT AAA AAA CCA CGA TCT 61 TCA TGT GGA AAC TGG AAT ATT TGC TGC GCA TAT GCT TGT GGA AAT ACA TAA CGA TGG TCC 121 GGT GAC TTT GTT ACT TGA TTC AAG AAA AGG TAT TTT GAA GTC ATC TTT GCT GTC TCT AGG 181 AGG ACT ATA TGC CTG AAT ACT CGC ATA GCA ATA AAA ACA ACT TTT TTG CCG AAA ACG ATG 240 241 TGA AGA ATT GTC ATC TAC TGC ATG TAT GTT GTG CAC CCG ATT TGG CAA TTT CTT ATT TGT 301 CCG GTG CAC GTG GTG ATA TTT TCT TTT ACA ATC CTA ACA TAC ATC CAA AAG CTG AAT ACG 361 AGA AAC GAC ACG CCG AAG TGA TTA AAA TTG CTG CAC TCT TTA AAA TGA ATG TTC TGA AAG 421 TTC CTT ATA ATC CTG ACC TGT TCT TCA AGC TTA CTA AAG GAT TAA AAA ATG AAC CTG AAG 481 GCG GGA CAA GGT GCG AGA TTT GTA TAA GAA TGC GAC TAG AAA AAA CAA TGG AAT ACG CGA 541 AAG AAA ATG GCT ACA AGA GTG TTT CCA CAA CGC TAA CAG CCT CTC CAA AGA AAA ATG TAG 601 CGA TGA TTG TGA AGA TAG GAA AAG AAC TGG AAA AAA AAT ACG GTG TGG AAT TTT TGC CTA 661 ATG TGT ACC GCA AAA GTC CGC TTT ACA ACG ATG CGC AAA AGC TTA TAA CGA AAA TGG GTT 720 721 ATT TAC AGA CAA AAC TAC TGT GGT TGT ATT TTC TCA ATA AGA ACT TCC GTT ATA GTA GCC 780 781 ACT CAA GAA ACT AAA ACC GTA AAA AGT GGG GTC GAA GTA TGA AAA TAT ACC ACA AAT TAG 841 AAG AAG TTG AAG AAC ATA AGC GGT CGT ATG CAT CAA TTG CTT TTT CAT CGA AAG TCA GGG 901 TTG AAT ATG AAC ATG CTG GCG AAA AAC TTG CCC TCA TCC CTG TAA CTA TTG GAG ACC TTA 960 961 CGG TGG TTA TCG AAA TTG ACG ATG ATA GAG AAG TAT TCA ATA CTT TGT TGA ACG AGC ACA 1020 1021 TCA AAA ACT CTA TCC TGA AAC AGT TTC CGT ATC CGG AAG AGA TTA GAG GGT TAG CCA GAC

1081 ATT TTC GCA CAG AAT TGA AGA ATT TCA GAA TCT TGG TTG TAA AAT ACA ATA GTG TCG AAG

1141 1200	AAA AGG	TAA	TCT	CAA	GGT	ATT	CAC	TGT	CTA	ATA	TAA	CAT	TCG	GTG	TGG	TGT	CAT	ACA	ATA
1201 1260	AAT TTG	ATG	TCC	ATT	TGT	TAC	CAA	GTA	ATG	TAA	AAG	TCA	GAC	CGA	AGC	CAG	GAT	ACT	GTC
1261 1320	TTT CAC	ATG	TTG	TCC	AAA	AGC	CTG	AAG	AAG	GTA	TCA	GGC	AAG	CAT	TCT	TGT	TAG	CCC	GGT
1321 1380	GGT TTG	GTG	GTG	GAA	GCT	ACG	ACC	AAC	TGC	CCA	TAA	TAG	CGC	TTG	AAA	GCA	CTG	ACA	TTG
1381 1440	ACC TTG	GAA	AGT	GGA	CAA	ATA	TAG	TCA	TAA	ACA	TCG	TTC	TGT	CAG	ATT	TTG	AAA	AGA	GGT
1441 1500	ATT TTT	CTG	GTA	TAA	TAA	AAA	AGC	TAA	ACG	AAT	TTA	GAA	GCG	AGA	CAT	ATT	TTG	ACC	CAT
1501 1560	TTG CTA	GGC	TTG	AAA	TGA	TAT	CAC	TTG	GCA	TAA	TAC	TCG	CCA	AGT	CAG	AGG	GAG	GAG	GTA
1561 1620	ACT TTG	AAC	CAG	ACA	GTT	ACG	ATA	TCA	TTT	AGA	gca	CTT	ACT	gaa	AAT	ATA	AAA	TTA	GCA
1621 1680	CGA GTT	GTT	ATA	CAT	act	TTT	CTA	ACA	TTC	CGA	GGA	GTG	TTC	gat	AAA	GAT	АТА	TTC	GAT
1681 1740	ACG GAA	TTG	GC T	GTA	AAC	GAA	GCG	ATT	GCA	AAC	ATT	ATT	CAG	CAT	ACA	TAC	AAA	GGT	GAA
1741 1800	CCA AAC	TAC	GTT	GTG	ATG	ACG	стс	AAT	TGG	ATA	gaa	CCA	GAT	ACA	CTC	GAA	GTG	TTA	CTC
1801 1860	CGC GAT	TTT	GGT	CCA	AAA	GTG	GAC	CCA	ACG	AAA	ATC	AAA	CCA	CGA	GAT	TTA	GAT	GAT	ATC
1861 1920	AGA CCA	GGA	GGA	CTC	GGA	GTT	TAT	ATA	ATT	CAA	CGC	ATC	TTC	GAC	ATT	ATG	GAA	TTC	CGA
1921 1980	AAC GTG	AGT	CAT	GGA	AAT	TTA	CTT	TAT	CTA	AAA	CGC	TCC	TTC	TTA	ATA	CCT	ccī	AAA	AAG
1981 2040	CAG GAG	CTT	GGG	AAT	TTA	AAT	AAT	GAA	ccc	TAT		GAA	TAT	TGA	AAA	AAC	CGT	CAA	AAC
2041 2100	GGG GGA	AAG	AAG	ACA	AAT	GGG	CTT	GCT	CAC	AGG			AAA	AAA	TCC	ATC	TTT	CAT	GTC
2101 2160	TGC ATT	TTT	TGG	CTT	TIT	GGC	AGC	ACA	ATT	TTT	GAA	AGT	GGT	GAT	ATA	CAA	AGA	TTT	CCG
2161 2220	CGT ATT	TGG	TAG	ATA	CGG	TGG	TAT	GCC	CAG	TGC	TCA	TGT	TGC	AAC	AAC	CTC	AGC	ATT	AGC
2221 2280	TTG GGC	TGT	TGG	TTA	CAC	TAC	AGG	TIT	TGA	TTC	ACC	GCT	TAC	AGC	CAT	CGC	TGC	AAT	TTT
2281 2340	CCT TGC	TAT	TAC	AAC	AGC	TGA	TGC	TGT	TGG	TTT	ACG	AAG	AAA	TGT	CGA	ccc	CAA	TAA	AGG

2341 2400	ACA TAC	ACT	AAT	GGA	AGC	TAT	CTA	TGG	CTT	CTT	ACT	TGG	GTG	gat	AGT	CGC	TCT	GCT	TAC
2401 2460	GGT TAA	GTT	GTA	TCG	ATA	ATT	TTG	TAA	GAG	TTG	TAG	TGA	AAT	AGC	CCA	AGT	CTT	TTT	TCG
2461 2520	CAA TTA	CAT	CAT	AAT	GCC	AGG	AGG	GTA	ATT	TAC	AAT	GTT	TTT	TAG	ATT	ACC	ATT	TAA	AGT
2521 2580	TTT TGT	TTT	TGC	AGT	TTT	GTT	GCT	TGC	CAT	CTC	GTT	AAC	AAG	TGT	TGT	TAG	TTT	TGG	ACA
2581 2640	AGA TGA	TGA	GCA	GAT	AAA	AAC	ACC	AAA	TTG	GTT	TAG	AAG	TGC	GGT	GAT	TAA	GAA	AAG	AGC
2641 2700	TGG TAT	GAA	TCT	AAA	GAC	CGC	ccc	AGA	GTT	TGT	AGA	TGA	CCT	ATG	GAA	TGC	GAT	ATA	CAC
2701 2760	TAT AGG	CAC	AAA	ATA	CAA	CGT	TCC	ccc	AAC	GCT	TAT	AGC	CGC	TGT	CAT	TTC	TGT	AGA	AAG
2761 2820	CAA CTT	CGC	CAA	CGT	gaa	AGG	TGC	TGG	AGA	CGT	GGT	AGG	AAT	GAT	GCA	AAT	TTC	TAT	CTC
2821 2880	CAC AGC	CAA	AAA	TAT	ATC	gaa	ACT	CCT	CGG	CCT	CGA	ACA	ACC	AAA	AA A	CGG	TTG	GGA	TGA
2881 2940	GCT CCT	CAC	AAA	TTA	TTG	GTT	GAA	TAT	AAC	TTA	CGG	TAC	CGC	ATA	CAT	CGC	TTA	TCT	TTA
2941 3000	CAA AAA	GCA	TGG	AAC	TTT	ACA	GAA	AGC	GCT	CGA	AGA	ATA	CAA	CAA	CGG	AAA	AAA	TAA	AAC
3001 3060	TAA ATA	CGC	CCA	GCT	GAT	ACT	ACA	ACA	ATA	CAA	CCT	ATA	CGA	GAG	CCT	CCA	TTC	TGC	TGA
3061 3120	AAT AAG	AAA	TAA	CCA	GCA	ATT	GGA	TAC	AGA	TAA	TTC	TTC	GAC	ATC	TTC	TGA	AGC	AAC	AGA
3121 3180	TAC TTT	GAA	TAC	AAC	CAG	TGC	AAC	AAA	TTC	ACA	ACC	AAC	ATC	AGA	TGC	ATC	AAA	TAC	ATC
3181 3240	AGT TAA	CAC	TTC	AGA	AAT	CAA	GTT	ccc	GCC	TCT	TTT	CGG	AGT	TGC	AGG	TTA	TTA	AGA	TAT
3241 3300	TTG TTC	GGT	AGT	TAC	TTA	GGA	ATG	TGG	GGT	GTA	TAG	TTT	GGA	AGA	TGA	AAA	AAT	GAA	ACC
330 <u>1</u>	TGA AAC	GAT	AGT	AAA	AAT	TGA	ACA	TTT	ATC	TTT	TTC	TTA	ccc	GAG	TTT	CAG	TCT	CAA	AGA
3361 3420	TGT AAG	TTT	TGA	GGT	TCG	GAA	GGG	AAG	TTT	CTT	CGG	CAT	TAT	TGG	ACC	AAA	TGG	TTC	GGG
3421 3480	AAA AAC	CAC	GCT	ACT	CTC	ACT	CAT	TAT	GAA	ATT	CCA	AAA	GCC	AAA	AAG	TGG	GAA	AAT	AAC
3481																			

3541 CAT CGC TCA AGA CTT TAA CCC TAC ATA CGA TTT CAC AGT TGA AGA ATT GGT CGA AAT GGG

- 3601 AGG AAT CCC CCG CTC ACC ACA TIT TIT CGA AAC ACC TGT TTA CGA GGA AGA ATT AGA AAA 3660
- 3661 TGC ACT CAA AAC TGT TGA TTT GCT TGA ATA CCG AAA AAG AAT ATT CTC CAC TCT TAG TGG
- 3721 AGG ACA ACA GCG CAG GGT CTT GAT TGC ACG CGC AAT CTA TCA AAA CAC ACC TAT CAT CAT 3780
- 3781 TGC TGA TGA ATT GGT TAA TCA CTT GGA TTT AGG GCA AGC AAT TAA AGT GTT AGA TTA TCT 3840
 - 3841 AAA ACA ACT TAC CGA ATG TGG AAA GAC GAT AAT TGG ACA TTC CAC CTG CAG CCC GG 3896

Archaeoglobus lithotrophicus TF2 (5phl)

SEQ ID NO:40

1 ATG TGC TGC AAG GCG ATT AAG TTG GTA ACG CCA GGT TTT CCC AGT CAC GAC GTT GTA AAA 61 CGA CGG CCA GTG AAT TGT AAT ACG ACT CAC TAT AGG GCG AAT TGG GTA CCG GGC CCC CCC 121 TCG AGG TCG ACG GTA TCG ATA AGC TTG ATA TCG AAT TCC GTA CGA AAT GCG GGA AAG AGA 181 GAA GGA AAA GGA AAG AGA GCA CAG ATT TGG AAA TGA GAC AGA ACA CGA GGA AGA GCA TGG 241 TAT GGC AGA GCG TGA AAG AGC ACA TGA GAA CGA GTC TGA AGA AAT GGG CAA GGG CGT TGG 301 CAT GGG CGC CCA TGG AAT GAA GAT GGG CAA AGA AGC TCG CGA AAT GGT GAA GGA AGA ATA 361 CAA GGA AGC AAA GGA GAG ATA CAA GAA GGC TAG AGA AGA GTT TGA AAG AGC AAA GAA GAT 421 GGG ATT GGA CAT CAG AGA GGA GCG CGG ATT CAA GAT GGC CAA GGG ATT CAT GGT AGC TGG 481 ACT AGA CGT TGC TGA GAT GTG GCT GGA GAG ACT GAA GGT ACA GGT CAT GAA TAT GGG TGA 541 AGA GGC CAA GAT CAC AGA GGA GAC CAA ACT GGA GCT GCT CGC AAA GAT CGA CGA GAA GCT 601 TGC AGA AAT CAA AGA GCT GAA GAA CGA AAT CAA TGA GAC CTC CTC ACC TGA AGA GCT GAT 661 AGA AAC TGT CAA GAA AAT CAG AAA GGA GTG GAG AGA AAT CAG AGA TGA AAT GAG GGC TCT 721 TAC TGG CTA TGT CGC CGT TGC CAA GGT GGA AAA GCT TGT TGA AAA GGC CAA GCA GGT AGA 780 781 GCT AAT GCT TGA GGC AAA GAT CGA GGA GCT CGA TGC TGC AGG AGT TGA TAC AAC CAA ACT 841 CGA GGC AAC ACT CGA GGA CTT CTC GGC AAA GGT TAA TGA AGC AGA AGA TTT GAT TGA CAA 901 GGC TGA AAA TCT GTT CGA GGA AGG CAA CAT TGC TGA AGG ACA CAT GAC TCT CAA GGA AGC 960 961 CAT AAA GAC TCT CAA GGA AGC CTT CAA GGA TGT CAA GGA AGT TGT CAG CGA GAT GAA GGA 1021 AAT GAA CCA GTA TAG AGT TAG GGA GGG CAA GAT CTT CTA CGG AAA CGA GAC TGG AGA AGT 1081 CTG GGT GGA TGG TAA TGG TAC TGC TGA GTT TAA CGG TAC CGG TAT CGT TGT GAT CAG AGG

1141 AAA CGC AAC ACT TGA GGT CGC ACC AGA AGA TGC GAT CGT GAC ACT GGT CGG CTT CGG CGT - 1200

- 1201 GAA GAG CGT TGA GGG TGG CGT TTC AAG AGT CAG CGG AGA AGG TAA GGC AGT AAT CAG AGG
- 1261 AGA AAA CCT CAC CGT CAA GGT GGA AGG TGA CGA CTT CAA GCT CAT AGT GAA GGG CTA CGG
- 1321 TAC ACT CAA ACT CGA TGG TGA GGG TGA ATA CAG GGT AAA GAA GAG CCC ACA GGA AGA GAT
- 1381 GAC ATT TAA ACT CTT TCT TCA ACT CTA GCA GTT TGA GCA TTG CAT TTC CAA GAT TTT TGC
- 1441 TGT TAG CTT CGG GAC AAC TTT GAA AAT ACG TCG AGA CAG GCT CAA ATG TTG TCC CAG CAT 1500
- 1501 TGC AGC TTT CGG CAA AGC GAA CGA GAT TTG CGT TCC GCT CCC CAG CCC AAC ATG GCT TCT 1560
- 1561 GTA ATC TGA AAA AAC TTC AAG TTC AAC AGC TTT CCC AAA AAC ATC CAA AAG CTT TTC CGC 1620
- 1621 AAC ACT TCT AAA TCT TTC GAG ATT TAT TGC ATT TCC TTT CAC CGA AAT GCT ATC GGA TTC 1680
- 1681 TOT TOO CAC AAC CTC GAT ATG CGG CTC TTC CAG AGC AAT ACC CAC TCC ACC GTC AAT CCT 1740
- 1741 TCC AAC CTG GCC GTT CAA ATC AAT GAG CGT GAT ATG AAT TCT CGA CGG AGT TTT AAC CTT
- 1801 AAC ATA CAT CTA TAG AAT TTA AAC GGT AAT TAC TTA AGA AGT TTT GGT TTT GCG AAA AAG
- 1861 AGT TCA AAA TTC ATT CTT TTA ACT GCA CTA CAG CTC ATC TGT GCC TTT TCT CCT TAA TTC
- 1921 GAT TIT TOT GAG ATA GTT CTG GTA TOT CGT ATC AAC TAT GTA AGC CTC GGG AGC TAT TAC 1980
- 1981 AGG CAG ATG ATA ACC GGT GAA TAT CCT TAT TAT CTC TCC AGC CTG AAC CGA GCA TGT CAG
- 2041 TGC ATA TGA TAT CGG ATC GTG ATC GAT GTG AGG ATA CTC CAC CTC GAA GAA AGA CAC ACC 2100
- 2101 ATC AGG CAG GAA AGT AGT AAT TAT ATC GGG AAT AAA TGG AGC TCC GAG CTC TTC AGC AAC 2160
- 2161 TTT TGC AGC CAT TGA AAT GTG CTT ATG AGC AAC AAC AAC AAT ACC TTT CAA CTG TCT 2220
- 2221 CCT GAG TTC TTT ATA ATC ATG CGG GAA GGG ATA AGA GAT TAT ACA CGA ATC AGA ACT CAT 2280
- 2281 AGG ATG CAC AAC ATC ATA ATC GTT TGC CTC AAG TGG CTT TAT GCT GGC ATC AAG CCT CAC

2341 ATC CAT TGG TGT AAC TAC ATC TCC AAT ATA CCG AAT GCA ACC AAC ACC ACT TCT CCA GAG -

- 2401 CAA TTC CAT GAG CAT TCT GCT TCC GAT GAC AGC GAC ACT AAA GTT CCT GAG ATA ATC TAT 2460
- 2461 CTT TTC TTC ATC TGC CAT CCC ATA CCA GGA AAT TTT TCT CAT GGC AAT AGC CCC GCA TCC
- 2521 ATT AAA TGG TAT TAA TTT TTT GCC GTA TTT TGA GGA GGT AGA TAT TAA CCA ATT ATT TTC
- 2581 AAA CCA TTT AAG GGC ATC GAT GAA ACA TCC CAA AAC CAG TTC AGC AAA AAA TTA AAT CAC
- 2641 TGC CAC ACA TTG AGG ACC CCA AAA TGG TGT GAG AAA TGG ACG AAC TGG GAG GAG TTA TTT 2700
- 2701 TTG ATC TGA TAG AAG AGG AGC CCG AAG TTG AGG AGG ACG ACG AGA TTA AGC TCG CAG AGA
- 2761 TAT ACA GGC TTG CTA CAA AAC TTA TAA AGT TAC TCG AAG ATC TCA AAA GCC ATG AGC TTA 2820
- 2821 AAG AGT CAG CAT CTC TTA TGC TCA TAA AGG AAA TTA TCG GTG AAG ACA GAG TTC TGG TTG
- 2881 GTT TAG CAT CAA AAA TGC TCC AGG ATA TGA GTC TCG GGT TCG AAG AGG ACG AAA AGT ACG 2940
- 2941 TTT CTT GAT TTT TGA ACT GTA TTT TCT ACA TGC TCT TTT CCC AAC CAC ATT CAG TTG CAT
- 3001 GCC ATA CGA AAA TTC CAA TGC CCA AAT CCT GGT AAA TGT ACT TTT TCA TAG TAA ATG CTG
- 3061 CCA AAC CCA GAT TAA ACT CAA TTT CAT CAA CAG GAA AAA GAA AGA ACG AAA AAA AGA CCT
- 3121 ACA ACA GTC CTA TAA TTG ACC AAA CTT GAT AGA TTA CAA ACA CCA CAG TTG GAA TCA AAG
 - 3181 CAC AGA TGA AAG CTT TCC GGA TTC CTG CAG CC 3212

Methanococcus thermolithoautotrophicus SN1 (14ph1)

Nucleic acid-SEQ ID NO:41 Amino acid-SEQ ID NO:42

1 ATG GAA ATA ATA AAC AAA TTT CTA AAA AAA ATT GGA TAT AAG AAA GAT GGA GAA AAA 60 1 Met Glu Ile Ile Asn Lys Phe Leu Lys Lys Ile Gly Tyr Lys Lys Asp Gly Glu Glu Lys 20 61 AAG GAC AAA TOT AAA ACC AAA ATA AAA ATT GAA GAA GAA AAA ACC ATG GAT ATC GAA ATT 21 Lys Asp Lys Ser Lys Thr Lys Ile Lys Ile Glu Glu Lys Thr Met Asp Ile Glu Ile 40 121 CCA AAA ATT GAA CCT ACT GAA AAT TTT AAT CGT GAT GAA ATT GTT TTT GAG GAA GAT AAT 180 41 Pro Lys Ile Glu Pro Thr Glu Asn Phe Asn Arg Asp Glu Ile Val Phe Glu Glu Asp Asn 181 GCC TAC GGT ATA TCC CAC AAA GGA AAT AGA ACA AAC AAC GAA GAC AAT ATT TTA ATT AGA 240 61 Ala Tyr Gly Ile Ser His Lys Gly Asn Arg Thr Asn Asn Glu Asp Asn Ile Leu Ile Arg 241 AAA ATA AAA GAT ACC TAC ATA TTA GCA GTT GCA GAT GGT GTC GGA GGG CAC AGC TCA GGA 300 81 Lys Ile Lys Asp Thr Tyr Ile Leu Ala Val Ala Asp Gly Val Gly Gly His Ser Ser Gly 301 GAT GTT GCA TCA AAG ATG GCA GTG GAT ATT TTA GAA AAC ATT ATC ATG GAA AAA TAC AAT 360 101 Asp Val Ala Ser Lys Met Ala Val Asp Ile Leu Glu Asn Ile Ile Met Glu-Lys Tyr Asn 361 GAA AAC CTA TCA ATT GAA GAG ATA AAA GAA CTT TTA AAA GAT GCA TAC ATT ACG GCA CAC 420 121 Glu Asn Leu Ser Ile Glu Glu Ile Lys Glu Leu Leu Lys Asp Ala Tyr Ile Thr Ala His 421 AAC AAA ATA AAA GAA AAC GCT ATT GGA GAT AAA GAG GGA ATG GGA ACA ACA CTA ACA ACT 480 141 Asn Lys Ile Lys Glu Asn Ala Ile Gly Asp Lys Glu Gly Met Gly Thr Thr Leu Thr Thr 481 GCA ATA GTT AAA GGG GAT AAA TGC GTT ATA GCA AAC TGC GGG GAT AGT AGG GCT TAT TTA 540 161 Ala Ile Val Lys Gly Asp Lys Cys Val Ile Ala Asn Cys Gly Asp Ser Arg Ala Tyr Leu 541 ATT AGA GAT GGA GAA ATA GTT TTT AGA ACA AAA GAC CAC TCT TTG GTT CAG GTT TTA GTA 181 Ile Arg Asp Gly Glu Ile Val Phe Arg Thr Lys Asp His Ser Leu Val Gln Val Leu Val 200

	601	GAT	GAA	GGA	CAT	ATT	TCA	GAG	GAG	GAC	GCA	AGG	CAT	CAT	CCA	ATG	AAA	AAT	ATC	ATT	ACC
	201	Asp	Glu	Gly	His	Ile	Ser	Glu	Glu	Asp	Ala	Arg	His	His	Pro	Met	Lys	Asn	Ile	Ile	Thr
220																					
	661	TCA	GÇA	TTG	GGA	TTG	GAT	GAA	TTT	AAG	GTA	GAT	GAT	TAC	GAA	TGG	GAT	TTA	ATT	GAT	GGT
	221	Ser	Ala	Leu	Gly	Leu	Asp	Glu	Phe	Lys	Val	Asp	Asp	Tyr	Glu	Trp	Asp	Leu	Ile	Asp	ĠΊγ
240	'																				
	721	GAT	GTA	TTA	TTG	ATG	AGC	TCC	GAT	GGG	CTT	CAT	GAT	TAT	GTC	AGT	AAG	GAA	GAT	ATT	TTA
	241	Asp	Val	Leu	Leu	Met	Ser	Ser	Asp	Gly	Leu	His	Asp	Tyr	Val	Ser	Lys	Glu	Asp	Ile	Leu
260)																				
	781	AAA	ACT	GTA	AAA	AAT	AAT	GAT	CAC	CCA	AAA	GAT	ATT	GTA	GAT	GAA	TTA	TTC	AAT	ACT	GCA
840	261	Lys	Thr	Val	Lys	Asn	Asn	Asp	Hıs	Pro	Lys	Asp	Ile	Val	Asp	Glu	Leu	Phe	Asn	Thr	Ala
280)																				
	841 281	TTA Leu	AAA Lys	GAG Glu	ACA Thr	AGG Arg	GAC Asp	AAT Asn	GTG Val	AGT Ser	ATT Ile	ATT Ile	CGT Arg	ATA Ile	879 293						

-90**-**

Pyrolobus fumarius 1A (1ph1)

SEQ ID NO 44-amino acid

1 ATG ACT CTG CTA GCC CTG TAT CAG AAT AAA CGT GTT ATC GTC AAG CTT GGC TGG GGG AGC 60 1 Met Thr Len Leu Ala Leu Tvr Gln Asn Lvs Arg Val Ile Val Lvs Leu Glv Trp Glv Ser 61 GGC ACT AGC CAA ATA ACT AAC GAG GCG CAA GTG CTG AGC GTA TTG CAC GAT ATG CCT ATA 120 21 Gly Thr Ser Gln Ile Thr Asn Glu Ala Gln Val Leu Ser Val Leu His Asp Met Pro Ile 121 GTG CCC AGA CTG CAT ACC CGT CTA GAC TTA GAT GAT GTC AAG CTC GTT GCG ATA GAG TAC 41 Val Pro Arg Leu His Thr Arg Leu Asp Leu Asp Val Lys Leu Val Ala Ile Glu Tyr 181 ATA CCC TAC AAG AGC CTT AAC GCC GTC GGC CGC TTG AAC CCC CTT AAG GCT GTC ACA GCC 240 61 Ile Pro Tyr Lys Ser Leu Asn Ala Val Gly Arg Leu Asn Pro Leu Lys Ala Val Thr Ala 241 GTC TTC TAT ACA CTC GCA TCG CTA GTC CAT ATC CAC GGC CGT GGT TTT GCT CAT TGC GAC 81 Val Phe Tyr Thr Leu Ala Ser Leu Val His Ile His Gly Arg Gly Phe Ala His Cys Asp 301 CTA AAG CCG GGT AAC GTT ATA CCA GTT CCC AAG CGT GGC ATG GTG TTC ATC GAC TTT GGT 360 101 Leu Lys Pro Gly Asn Val Ile Pro Val Pro Lys Arg Gly Met Val Phe Ile Asp Phe Gly 120 361 GTT GCA CGA CCT TTT GAC GCT GCG GGC TTC GCG GCA GGA ACA CCA GGG TAT ACG TGC CCA 121 Val Ala Arg Pro Phe Asp Ala Ala Gly Phe Ala Ala Gly Thr Pro Gly Tyr Thr Cys Pro 140 421 GAG GCT CTC GGC GGC GAG ACC CCC GGC TCT GGC TGC GAT CTC TAC AGC CTT GCC GGC ATA 480 141 Glu Ala Leu Gly Gly Glu Thr Pro Gly Ser Gly Cys Asp Leu Tyr Ser Leu Ala Gly Ile 481 TAC TAC TAC TTG GTT ACC GGG TTA AGC CCG CCA CGC GAC CCA AAA GAG TTC GCC AAG GCG 540 161 Tyr Tyr Tyr Leu Val Thr Gly Leu Ser Pro Pro Arg Asp Pro Lys Glu Phe Ala Lys Ala 541 CTC TCG TTG GCT CCC GCT CCA AGT AGC CTC TTG GAA CTG TTC ACA CAG CTG GTG CTG GAT 600 181 Leu Ser Leu Ala Pro Ala Pro Ser Ser Leu Leu Glu Leu Phe Thr Gln Leu Val Leu Asp 601 CCC GAG TAT CGT AAC AGC CTT GAT CCT CTC CAG CTG TTG AAG ATT GTT GCA TCT TTT AAC 660 201 Pro Glu Tyr Arg Asn Ser Leu Asp Pro Leu Gln Leu Leu Lys Ile Val Ala Ser Phe Asn 661 CCG CAA CTG CTA GTC CCT CAT ATC GTT ATA GAT GGT GTT TAC AAG CCG CTA GGT TAC GGC 221 Pro Gln Leu Leu Val Pro His Ile Val Ile Asp Gly Val Tyr Lys Pro Leu Gly Tyr Gly 721 GAG GTA AGC ATA GGC TCT AGA GGC GTT ATA CGT GTT GAT GGA CGA CCA GTG TAC CTC GCG 780 241 Glu Val Ser Ile Gly Ser Arg Gly Val Ile Arg Val Asp Gly Arg Pro Val Tyr Leu Ala 260 781 GTT AAG AGG CAT GTG AGG GGC ACA AGT ATG TAC GCG TAT ACG GAT CTT GTC GTG TTT AGG 261 Val Lys Arg His Val Arg Gly Thr Ser Met Tyr Ala Tyr Thr Asp Leu Val Val Phe Arg 280

841 AGA GGC GAG AAA CTC ATA GTG AGA AGC GGT GAG AGT ATA GAC CTA GAG TTT AAC GAC CTG -281 Arg Gly Glu Lys Leu Ile Val Arg Ser Gly Glu Ser Ile Asp Leu Glu Phe Asn Asp Leu 300

901 GTG TTG GAC AAC CAC ATA CTA TAC GTA TTT ATC CTT CCG GAA AGG CCC 951 301 Val Leu Phe Asp Asn His Ile Leu Tyr Val Phe Ile Leu Pro Glu Arg Pro 317

-92-

Thermococcus celer (25ph2)

SEQ ID NO:46-amino acid

1 ATG GAC ATC AGG GCC GTT GTT TTT GAC CTC GAC GGG ACG CTT GTG GGT GCT GAG AAG ACT 1 Met Asp Ile Arg Ala Val Val Phe Asp Leu Asp Gly Thr Leu Val Gly Ala Glu Lys Thr 2.0 61 TTC AGC GAG ATA AAG TCC GAG CTT AAA GAA CGG CTG ATT TCC TTA GGG ATT CCC AGG GAG 21 Phe Ser Glu Ile Lys Ser Glu Leu Lys Glu Arg Leu Ile Ser Leu Gly Ile Pro Arg Glu 40 121 CTC GTT GGA GAG CTA ACG CCG ATG TAT GAG GGC CTT ATC GAG CTG TCC AGA AAA ACG GGC 41 Leu Val Gly Glu Leu Thr Pro Met Tyr Glu Gly Leu Ile Glu Leu Ser Arg Lys Thr Gly 181 AGA CCT TTC GAA GAG ATG TAC TCA ATT CTC GTC AAT CTT GAA GTT GAA AGG ATA AGG GAC 61 Arg Pro Phe Glu Glu Met Tyr Ser Ile Leu Val Asn Leu Glu Val Glu Arg Ile Arg Asp 241 AGC TTT CTC TTC GAG GGG GCA AGG GAG CTC CTC GAC TTT CTT GTG GGG GAG GGA ATA AAG 300 81 Ser Phe Leu Phe Glu Gly Ala Arg Glu Leu Leu Asp Phe Leu Val Gly Glu Gly Ile Lys 301 CTT GCC CTC ATG ACC CGG AGC TCC AGA ATG GCT GCC CTT GAG GCC CTG GAG CTT CAC GGC 360 101 Leu Ala Leu Met Thr Arg Ser Ser Arg Met Ala Ala Leu Glu Ala Leu Glu Leu His Gly 361 ATT AAG GAC TAC TTT GAG ATT ATT TCA ACG AGG GAT GAT GTC CCC GAG GAG CTG AAA 420 121 Ile Lys Asp Tyr Phe Glu Ile Ile Ser Thr Arg Asp Asp Val Pro Pro Glu Glu Leu Lys 140 421 CCG AAT CCT GGC CAG CTG AGG AGA ATC CTC GGT GAG CTC AAC GTT CAA CCA GAG AAA GCC 141 Pro Asn Pro Gly Gln Leu Arg Arg Ile Leu Gly Glu Leu Asn Val Gln Pro Glu Lys Ala 481 ATC GTC GTT GGA GAC CAC GGC TAC GAT GTC ATC CCT GCC CGG GAG CTC GGC GCT CTG AGC 540 161 Ile Val Val Gly Asp His Gly Tyr Asp Val Ile Pro Ala Arg Glu Leu Gly Ala Leu Ser 180 541 GTC CTT GTC ACC GGC CAC GAG GCT GGC AGA ATG AGC TTT CAG GTT GAA GCC GAG CCA AAC 600 181 Val Leu Val Thr Gly His Glu Ala Gly Arg Met Ser Phe Gln Val Glu Ala Glu Pro Asn 601 TTT GAG GTC GAG AAC CTC ATT CAC CTC AGG AAG CTC TTC GAG AGG CTC CTG TCG AGC TAC 660 201 Phe Glu Val Glu Asn Leu Ile His Leu Arg Lys Leu Phe Glu Arg Leu Leu Ser Ser Tyr 661 GTT GTT GTT CCC GCT TAC AAC GAG GAG AAG ACC ATC AAG GGG GTA ATA GAG AAT CTT CTC 720 221 Val Val Val Pro Ala Tyr Asn Glu Glu Lys Thr Ile Lys Gly Val Ile Glu Asn Leu Leu 721 AGG TAT TTC AAA AAG GAC GAG ATA ATC GTC GTG AAC GAC GGC TCC AGG GAT AGA ACG GAG 780 241 Arg Tyr Phe Lys Lys Asp Glu Ile Ile Val Val Asn Asp Gly Ser Arg Asp Arg Thr Glu 781 GAG ATA GCT CGT TCT TAC GGA GTC CAC GTT CTT ACG CAT CTC GTC AAC AGG GGG CTT GGT 840 261 Glu Ile Ala Arg Ser Tyr Gly Val His Val Leu Thr His Leu Val Asn Arg Gly Leu Gly 280

841 GGG GCC CTC GGA ACG GGC TTT GCC TAT GCC ATC AGA AAA AAC GCC AAA CTT GTC CTC ACA 281 Gly Ala Leu Gly Thr Gly Phe Ala Tyr Ala Ile Arg Lys Asn Ala Lys Leu Val Leu Thr 901 TIT GAT GCC GAC GGC CAG CAC CIT ATA AGC GAC GCC CTC CGC GTC ATG AGG CCA GTT GCG 301 Phe Asp Ala Asp Gly Gln His Leu Ile Ser Asp Ala Leu Arg Val Met Arg Pro Val Ala 961 GAG GGC AGG GGC GAC TTT GCG GTC GGC TCA AGG CTC AAA GGT GAC ACG AGC CAG ATG CCC 321 Glu Gly Arg Ala Asp Phe Ala Val Gly Ser Arg Leu Lys Gly Asp Thr Ser Gln Met Pro 340 1021 CTC GTG AAG AAG TTC GGC AAC TTC GTT CTA GAT GCC GTG ACC GCG GTT TTT GCT GGT AAA 1080 341 Leu Val Lys Lys Phe Gly Asn Phe Val Leu Asp Ala Val Thr Ala Val Phe Ala Gly Lys 360 1081 TAC GTC AGC GAC AGT CAG AGC GGG TTA AGG TGT CTA AGC GGC GAC TGC CTG AGG AAA ATC 1140 361 Tyr Val Ser Asp Ser Gln Ser Gly Leu Arg Cys Leu Ser Gly Asp Cys Leu Arg Lys Ile 380 1141 AGG ATA ACC TGC GAC CGC TAT GCC GTG TCG AGT GAG ATT ATA ATA GAG GCC TCC AAA GCG 381 Arg Ile Thr Cys Asp Arg Tyr Ala Val Ser Ser Glu Ile Ile Ile Glu Ala Ser Lys Ala 1201 GGC TGT AGA ATT GTC GAA GTT CCT ATC AAG GCT GTT TAC ACT GAG TAC TTT ATG AAG AAG 401 Gly Cys Arg Ile Val Glu Val Pro Ile Lys Ala Val Tyr Thr Glu Tyr Phe Met Lys Lys 420 1261 GGG ACG AAC GTT TTA GAG GGC GTT AAG ATA GCC CTG AAC CTT CTC TTT GAC AAA CTG AGG 421 Gly Thr Asn Val Leu Glu Gly Val Lys Ile Ala Leu Asn Leu Leu Phe Asp Lys Leu Arg

Aquifex pyrophilus (28phl)

SEQ ID NO:47 and 48

1 ATG GAA AAT CTT GAA AAA CTC CTT GAA GTG GCA AAG ATG GCA GCC CTT GCC GGA GGA CAG 60 1 Met Glu Asn Leu Glu Lys Leu Leu Glu Val Ala Lys Met Ala Ala Leu Ala Gly Gly Gln 20 61 GTA TTA AAG GAA AAC TTC GGA AAG ATT AAG CTT GAA AAC ATT GAA GAA AAG GGA GAG AAG 21 Val Leu Lys Glu Asn Phe Gly Lys Ile Lys Leu Glu Asn Ile Glu Glu Lys Gly Glu Lys 40 121 GAC TTC GTG AGC TAC GTT GAT AAA ACC TCC GAA GAG AGA ATA AAA GAG CTA ATA CTT AAG 41 Asp Phe Val Ser Tyr Val Asp Lys Thr Ser Glu Glu Arg Ile Lys Glu Leu Ile Leu Lys 181 TTC TTT CCC GAC CAC GAG GTC GTG GGG GAA GAG GGA AAG GAG GGA AAA GAA AGC CCT 240 61 Phe Phe Pro Asp His Glu Val Val Gly Glu Glu Arg Gly Lys Glu Gly Lys Glu Ser Pro 80 241 TAC ARA TGG TTC ATA GAC CCC CTT GAT GGG ACC AAG AAC TAC ATA AAG GGC TTT CCC ATA 81 Tyr Lys Trp Phe Ile Asp Pro Leu Asp Gly Thr Lys Asn Tyr Ile Lys Gly Phe Pro Ile 301 TIT GCA GTC TCC GTC GGA CTC GTT AAG GAA AAC GAA CCT ATA GTG GGA GCG GTT TAC CTT 360 101 Phe Ala Val Ser Val Gly Leu Val Lys Glu Asn Glu Pro Ile Val Gly Ala Val Tyr Leu 361 CCT TAC TTT GAT ACC CTA TAC TGG GCT TCA AAG GGA AGG GGA GCC TAT AAA AAC GGG GAG 420 121 Pro Tyr Phe Asp Thr Leu Tyr Trp Ala Ser Lys Gly Arg Gly Ala Tyr Lys Asn Gly Glu 421 AGG ATA AGC GTA AAG GAA AGG GGG GAG CTC AAG CAC GCG GCG GTT GTT TAC GGA TTT CCA 480 141 Arg Ile Ser Val Lys Glu Arg Gly Glu Leu Lys His Ala Val Val Tyr Gly Phe Pro 481 TCA AGA AGC AGG AGG GAT ATA TCT CTT TAC CTG AAT GTG TTT AAA GAG GTC TTT TAC GAA 540 161 Ser Arg Ser Arg Arg Asp Ile Ser Leu Tyr Leu Asn Val Phe Lys Glu Val Phe Tyr Glu 180 541 GTA GGT TCC GTT AGG AGG CCC GGG GCC GCA GCG GTT GAT ATA TGC ATG CTT GCG GAG GGC 181 Val Gly Ser Val Arg Arg Pro Gly Ala Ala Ala Val Asp Ile Cys Met Leu Ala Glu Gly 200 600 601 ATA TTT GAC GGG ATG ATG GAG TTT GAG ATG AAG CCA TGG GAC ATA ACG GCG GGA CTC GTA 660 201 Ile Phe Asp Gly Met Met Glu Phe Glu Met Lys Pro Trp Asp Ile Thr Ala Gly Leu Val 661 ATA CTG AAG GAA GCT GGA GGA TTT TAC ACA CTG AAG GGA GAC CCC TTC GGC ATC TCG GAC 221 Ile Leu Lys Glu Ala Gly Gly Phe Tyr Thr Leu Lys Gly Asp Pro Phe Gly Ile Ser Asp 240 721 ATA ATA GCG GGA AAC AGG ATG CTC CAC GAC TTC ATT CTC AAG GTT GTG AAT AAA TAC ATG 780 241 Ile Ile Ala Gly Asn Arg Met Leu His Asp Phe Ile Leu Lys Val Val Asn Lys Tyr Met

781 AAT AAT GAA AGC ACG 795 261 Asn Asn Glu Ser Thr 265

Bacillus thermoleovorans (68FYS)

SEQ ID NO:49 and 50

60	ATG	AGT	GAA	CAG	CCG	GTA	TTG	TCT	GTT	CAA	GGA	TTA	AGC	GGC	GGG	TAT	AGC	ATG	AAC	CGA
20	Met	Ser	Glu	Gln	Pro	Val	Leu	Ser	Val	Gln	Gly	Leu	Ser	Gly	Gly	Tyr	Ser	Met	Asn	Arg
61 120	CCG	GTT	CTG	CAT	GAC	GTA	ACC	TTT	CAG	GTT	GAA	CCG	GGT	GAG	ATG	GTG	GGT	TTG	ATC	GGC
40	Pro	Val	Leu	His	Asp	Val	Thr	Phe	Gln	Val	Glu	Pro	Gly	Glu	Met	Val	Gly	Leu	Ile	Gly
121 180	CTG	AAC	GGT	GCG	GGC	AAG	AGT	ACC	ACG	ATG	AAG	CAT	ATT	CTC	GGG	CTG	ATG	AAT	CCG	CAA
60 41	Leu	Asn	Gly	Ala	Gly	Lys	Ser	Thr	Thr	Met	Lys	His	Ile	Leu	Gly	Leu	Met	Asn	Pro	Gln
181 240	AAA	GGG	AGC	ATT	CAG	GTT	CAA	GGA	AAG	AGC	CGG	ACA	GAG	CAT	TCG	GAA	GCC	TAT	CAC	GGC
61 80	Lys	Gly	Ser	Ile	Gln	Val	Gln	Gly	Lys	Ser	Arg	Thr	Glu	His	Ser	Glu	Ala	Tyr	Hís	Gly
241 300	GCC	TTG	GCG	TTT	GTT	ccc	GAA	TCC	CCG	CTG	CTG	TAT	GAG	GAG	ATG	ACA	GTA	CGA	GAG	CAT
100	Ala	Leu	Ala	Phe	Val	Pro	Glu	Ser	Pro	Leu	Leu	Tyr	Glu	Glu	Met	Thr	Val	Arg	Glu	Hıs
301 360	CTG	GAA	TTT	ACG	GCG	CGC	TCC	TAT	GGC	GTA	TCC	CGT	GAA	GAT	TAT	GAG	GCA	CGT	TCG	GAG
101 120	Leu	Glu	Phe	Thr	Ala	Arg	Ser	Tyr	Gly	Val	Ser	Arg	Glu	Asp	Tyr	Glu	Ala	Arg	Ser	Glu
361 420	CAG	CTG	TCG	AAG	ATG	TTC	CGT	ATG	GAA	GAG	AAG	ATG	GAC	AGC	CTG	TCC	ACG	CAT	TTG	TCC
121 140	Gln	Leu	Ser	Lys	Met	Phe	Arg	Met	Glu	Glu	Lys	Met	Asp	Ser	Leu	Ser	Thr	His	Leu	Ser
421 480	AAA	GGG	ATG	CGC	CAA	AAA	GTG	ATG	ATC	ATG	TGC	GCA	TTC	GTA	GCC	AGA	CCG	TCC	CTG	TAC
141 160	Lys	Gly	Mec	Arg	Gln	Lys	Val	Met	Ile	Met	Суз	Ala	Phe	Val	Ala	Arg	Pro	Ser	Leu	Tyr
481 540	ATC	ATT	GAC	GAG	ccc	TTT	CTT	GGG	CTT	GAT	CCG	CTT	GGG	ATA	CGC	TCG	CTG	CTT	GAC	TTC
161 180	Ile	Ile	Asp	Glu	Pro	Phe	Leu	Gly	Leu	qeA	Pro	Leu	GΙγ	Ile	Arg	Ser	Leu	Leu	Asp	Phe
541 181	-	CTG Leu		-									-						91 97	

Pyrococcus furiosus VC1 (7ph1)

SEQ ID NO:51 and 52

1 ATG AAG AAA ATA ACT ATT AGT AGT TTG CTT CTA CTT TTA CTT ATT TCT ACC AAT TTG AAT 1 Met Lys Lys Ile Thr Ile Ser Ser Leu Leu Leu Leu Leu Leu Ile Ser Thr Asn Leu Asn 20 61 CTC GCA TAC GAT TCC CAA GAG AGC GGT ATT AAA AAT ATA ATA ATC CTC ATT GGA GAC GGC 21 Leu Ala Tyr Asp Ser Gln Glu Ser Gly Ile Lys Asn fle Ile Ile Leu Ile Gly Asp Gly 4.0 121 ATG GGA ATG AGT CAT GTC CAG ATT ACA AAG CTT GTT TAT GGT CAT CTA AAC ATG GAA GAG 41 Met Gly Met Ser His Val Gln Ile Thr Lys Leu Val Tyr Gly His Leu Asn Met Glu Glu 181 TTC CCA ATT ATT GGA TTC GAA CTT ACT GAG TCA TTA AGT GGG GAA GTT ACG GAC TCC GCT 61 Phe Pro Ile Ile Gly Phe Glu Leu Thr Glu Ser Leu Ser Gly Glu Val Thr Asp Ser Ala 241 GCA GCA GGA ACT GCA ATA GCA ACT GGA GTC AAA ACA TAT AAT CGA ATG ATT TCA GTT ACT 81 Ala Ala Gly Thr Ala Ile Ala Thr Gly Val Lys Thr Tyr Asn Arg Met fle Ser Val Thr 301 AAC ATA ACT GGA AAA GTT ACA AAT CTA ACT TTG CTT GAA ATA GCC CAG GTA CTT GGA 101 Asn Ile Thr Gly Lys Val Thr Asn Leu Thr Thr Leu Leu Glu Ile Ala Gln Val Leu Gly 361 AAA TCA ACT GGA CTT GTG ACT ACT ACT AGA ATT ACA CAC GCA ACC CCT GCA GTA TTT GCT 420 121 Lys Ser Thr Gly Leu Val Thr Thr Thr Arg Ile Thr His Ala Thr Pro Ala Val Phe Ala 421 TCC CAC GTT CCT GAC AGA GAT ATG GAA GAG GAA ATA GCG AGA CAG CTC ATA GCT CAC CGG 141 Ser His Val Pro Asp Arg Asp Met Glu Glu Glu Ile Ala Arg Gln Leu Ile Ala His Arg 481 GTC AAC GTC CTA TTA GGT GGA GGG AGA AAG AAA TTT GAC GAG AAT ACC CTA AAA ATG GCA 540 161 Val Asn Val Leu Leu Gly Gly Gly Arg Lys Phe Asp Glu Asn Thr Leu Lys Met Ala 180 541 AAA GAA CAG GGA TAT AAT ATA GTC TTC ACG AAA GAA GAG CTC GAG AAA GCA GAG GGT GAG 181 Lys Glu Gln Gly Tyr Asn Ile Val Phe Thr Lys Glu Glu Leu Glu Lys Ala Glu Gly Glu 200 601 TTT ATT CTA GGG CTT TTT GCA GAT AGC CAC ATT CCT TAC GTA TTG GAC AGA AAA CCA GAA 660 201 Phe Ile Leu Gly Leu Phe Ala Asp Ser His Ile Pro Tyr Val Leu Asp Arg Lys Pro Glu 661 GAT GTT GGA CTT TTG GAA ATG ACT AAA AAA GCA ATT TCA ATA CTA GAG AAA AAT CCA AAT 720 221 Asp Val Gly Leu Leu Glu Met Thr Lys Lys Ala Ile Ser Ile Leu Glu Lys Asn Pro Asn 721 GGG TTC TTT CTC ATG ATT GAA GGG GGC AGA ATT GAT CAT GCA GCT CAT GAG AAT GAT ATA 780 241 Gly Phe Phe Leu Met Ile Glu Gly Gly Arg Ile Asp His Ala Ala His Glu Asn Asp Ile 781 GCA TCA GTT GTT GCA GAG ACT AAG GAG TTT GAT GAC GTT GTT GGA TAT GTT CTT GAG TAT 261 Ala Ser Val Val Ala Glu Thr Lys Glu Phe Asp Asp Val Val Gly Tyr Val Leu Glu Tyr 841 GCA AAA AAG AGG GGA GAT ACA CTA GTA ATA GTG CTG GCT GAC CAT GAG ACA GGG GGG CTT 900 281 Ala Lys Lys Arg Gly Asp Thr Leu Val Ile Val Leu Ala Asp His Glu Thr Gly Gly Leu

901	GGA	TTA	GGT	CTA	ACA	TAT	GGA	GAT	GCA	ATT	AAT	GAA	GAT	GTC	ATC	AGG	AAC	ATA	AAC	GCT
960 301 320	Gly	Leu	Gly	Leu	Thr	туг	Gly	Asp	Ala	Ile	Asn	Glu	Ąsp	Val	Ile	Arg	Asn	Ile	Asn	Ala
961 1020				AAA																
321 340	Ser	Val	Ser	Lys	Ile	Ala	Ser	Glu	Ile	Arg	Ala	Thr	Asn	Asp	Ile	Lys	Arg	Val	Ile	Lys
1021 1080				GGA																
341 360	Lys	Tyr	Thr	Gly	Phe	Glu	Leu	Thr	Glu	Asp	Glu	Ile	Asn	Tyr	Ile	Glu	Glu	Alā	Ile	Asn
1081 1140				GAA																
361 380	Leu	Ala	Asp	Glu	Tyr	Ala	Leu	Gln	Asn	Ala	Ile	Ala	Asp	Ile	Ile	Asn	Lys	Arg	Val	Gly
1141 1200				GTA																
381 400		_		Val																
1201 1260				AAT																
401 420	•			Asn																
1261 1320				TTT																
421 440				Phe														-		
1321 1380				ACC																
441 460	_			Thr																
1381 1440				GAA																
461 480			_	Glu													Asn	AST	Asn	GIÀ
1441 481	ATA Ile	ATC	GAG Glu	TTG Leu	GGA Gly	GAC Asp	GTG Val	CTC Leu	CTG Leu	ATT	CTA Leu	CAA Gln	GAG Glu	TCC Ser	148 494	2				

Pyrococcus furiosus VC1 (7ph2)

SEQ ID NO:53 and 54

1 ATG ATT AAC CAA ATA AAC TTC AAA ACC TCT CAT GGA GGA AGC AGA GAA GAA GGC TAC ATA 60 1 Met Ile Asn Gln Ile Asn Phe Lys Thr Ser His Gly Gly Ser Arg Glu Glu Gly Tyr Ile 20 61 AAC TTC TCG GCC TCT GTA AAT CCT TAT CCA CCA GAA TGG ACT GAT GAA ATG TTT GAG AGG 21 Asn Phe Ser Ala Ser Val Asn Pro Tyr Pro Pro Glu Trp Thr Asp Glu Met Phe Glu Arg 121 GCT AAA AAG ATA AGC ACC TTC TAT CCT TAC TAT GAA AAG CTT GAG GAA GAA CTC TCA GAT 41 Ala Lys Lys Ile Ser Thr Phe Tyr Pro Tyr Tyr Glu Lys Leu Glu Glu Glu Leu Ser Asp 60 181 CTA ATT GGG GAG CCA ATA ACT ATA ACT GGA GGA ATA ACA GAG GCA CTT TAC CTG CTT GGA 61 Leu Ile Gly Glu Pro Ile Thr Ile Thr Ala Gly Ile Thr Glu Ala Leu Tyr Leu Leu Gly 241 GTT TGG ATG AGG GGT CGG AAA GTA ATA ATC CCG AAG CAC ACC TAT GGG GAA TAC GAG AGG 300 81 Val Trp Met Arg Gly Arg Lys Val Ile Ile Pro Lys His Thr Tyr Gly Glu Tyr Glu Arg 301 ATC TCA CGC ATG TTC GGA GGT AGG GTG ATC AAA GGT CCC AAT GAC CCA GGA AAG TTA GCA 101 Ile Ser Arg Met Phe Gly Gly Arg Val Ile Lys Gly Pro Asn Asp Pro Gly Lys Leu Ala 361 GAA TIT GIT GAA AGA AAT TCA TTC GTG TTC TTC TGC AAT CCA AAC AAT CCA GAT GGA AAG 121 Glu Phe Val Glu Arg Asn Ser Phe Val Phe Phe Cys Asn Pro Asn Asn Pro Asp Gly Lys 421 TTC TAC CGA GAA AAA GAG ATG AAA CCT CTT TTA GAT GCC ATT CAA GAC ACT AAC TCA ATT 480 141 Phe Tyr Arg Glu Lys Glu Met Lys Pro Leu Leu Asp Ala Ile Gin Asp Thr Asn Ser Ile 481 TTG ATC TTG GAT GAA GCC TTC ATA GAC TTT GTT AAG AAA CCA GAA AGC CCA GAG GGA GAG 161 Leu Ile Leu Asp Glu Ala Phe Ile Asp Phe Val Lys Lys Pro Glu Ser Pro Glu Gly Glu 180 541 AAC ATA ATC AGG CTA AGG ACT TTT ACC AAA AGC TAC GGG CTC CCA GGG GTA AGG GTT GGA 181 Asn Ile Ile Arg Leu Arg Thr Phe Thr Lys Ser Tyr Gly Leu Pro Gly Val Arg Val Gly 601 TAT GTT ATT GGA TTT GTC GAT GCT TTC AGG AGC GTT AGA ATG CCA TGG TCA ATT GGC TCT 201 Tyr Val Ile Gly Phe Val Asp Ala Phe Arg Ser Val Arg Met Pro Trp Ser Ile Gly Ser 661 ACT GGG GTG GCC TTC TTA GAG TTC TTA CTC AAA GAT AAC TTC AAA CAC TTA AGA AAA ACC 221 Thr Gly Val Ala Phe Leu Glu Phe Leu Leu Lys Asp Asn Phe Lys His Leu Arg Lys Thr 240 721 CTC CCC CTA ATA TGG AAA GAA AAG GAG AGG ATT GAG AAA GAA TTG AAA GTT AAA AGC GAT 241 Leu Pro Leu Ile Trp Lys Glu Lys Glu Arg Ile Glu Lys Glu Leu Lys Val Lys Ser Asp 781 GCA AAT TTC TTC ATT ATG AAG GTC AGA GAA GGA ATA ATT GAA AAG CTA AAA GAG AAT GGC 840 261 Ala Asn Phe Phe Ile Met Lys Val Arg Glu Gly Ile Ile Glu Lys Leu Lys Glu Asn Gly 280 841 ATC CTT GTA AGG GAT TGC AAG AGC TTT GGA CTC CCT GGG TAC ATA AGG TTT TCA GTT AGA 281 Ile Leu Val Arg Asp Cys Lys Ser Phe Gly Leu Pro Gly Tyr Ile Arg Phe Ser Val Arg 300

901 AGG AGA GAA GAG AAT GAC AAA CTC ATA AAC ATC CTT AGA AAA ACA CTT AAT ACT 954 301 Arg Arg Glu Glu Asn Asp Lys Leu Ile Asn Ile Leu Arg Lys Thr Leu Asn Thr 318

What Is Claimed Is:

1. An isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide encoding an enzyme comprising an amino acid sequence selected from the group of amino acid sequences set forth in SEQ ID NOS:28-36;
- (b) a polynucleotide which is complementary to the polynucleotide of (a); and
- (c) a polynucleotide comprising at least 15 bases of the polynucleotide of (a) or (b).
- 2. An isolated polynucleotide selected from the group consisting of:
 - (a) SEQ ID NOS:19-27, 37-41, 43, 45, 47, 49, 51, or 53;
 - (b) SEQ ID NOS:19-27, 37-41, 43, 45, 47, 49, 51, or 53, where T can also be U; and
 - (c) fragments of a)or b)that are at least 15 bases in length and that will hybridize to DNA which encodes the amino acid sequence of any of SEQ ID Nos:28-36, 42, 44, 46, 48, 50, 52, or 54.
- 3. The polynucleotide of Claim 1 wherein the polynucleotide is DNA.
- 4. The polynucleotide of Claim 1 wherein the polynucleotide is RNA.

5. An isolated polynucleotide comprising a polynucleotide having at least 70% identity to a member selected from the group consisting of:

- (a) a polynucleotide encoding an enzyme encoded by the DNA contained in ATCC Deposit No. 97379, wherein said enzyme is selected from the group consisting of Ammonifex degensii KC4, Aquifex VF-5, M11TL, Methanococcus igneus KOL5, Thermococcus AED112RA, and Thermococcus celer, Thermococcus CL-2, and Thermococcus GU5L5.
- (b) a polynucleotide complementary to the polynucleotide of (a); and
- (c) a polynucleotide comprising at least 15 bases of the polynucleotide of (a) and (b).
- 6. A vector comprising the DNA of Claim 1 or Claim 2.
- 7. A host cell comprising the vector of Claim 6.
- 8. A process for producing a polypeptide comprising: expressing from the host cell of Claim 7 a polypeptide encoded by said DNA and isolating the polypeptide.
- 9. A process for producing a recombinant cell comprising: transforming or transfecting the cell with the vector of Claim 6 such that the cell expresses the polypeptide encoded by the DNA contained in the vector.

10. An enzyme of which at least a portion is coded for by a polynucleotide of claim 1, and which is selected from the group consisting of:

- (a) an enzyme comprising an amino acid sequence which is at least 70% identical to an amino acid sequence selected from the group of amino acid sequences set forth in SEQ ID NOS:28-36; and
- (b) an enzyme which comprises at least 30 amino acid residues to the enzyme of (a).
- 11. An enzyme of which at least a portion is coded for by a polynucleotide of claim 1, and which is selected from the group consisting of:
- (a) an enzyme comprising an amino acid sequence selected from the group of amino acid sequences set forth in in SEQ ID NOS:28-36, 42, 44, 46, 48, 50, 52, or 54; and
- (b) an enzyme which comprises at least 30 amino acid residues to the enzyme of (a).
- 12. A method for hydrolyzing phosphate bonds comprising:

administering an effective amount of an enyzme selected from the group consisting of an enzyme having the amino acid sequence selected from the group of amino acid sequences set forth in SEQ ID NOS:28-36, 42, 44, 46, 48, 50, 52, or 54.

FIGURE 1.

Ammonifex degensii KC4 Phosphatase(3AlA=3A2A) Complete gene sequence

1	ATGAGGGGGAGCGGAGTGCGGATACTTCTCACCAACGATGACGGCATCTTTGCCGAGGGT MetArgGlySerGlyValArgIleLeuLeuThrAsnAspAspGlyIlePheAlaGluGly
	CTGGGGGCTCTGCGCAAGATGCTGGAGCCCGTGGCTACCCTTTACGTGGTGGCTCCGGAC
21	LeuGlyAlaLeuArgLysMetLeuGluProValAlaThrLeuTyrValValAlaProAsp
	CGAGAGCGTAGCGCGGCCAGCCATGCTATCACCGTTCACCGCCCCCTGCGGGTGCGGGAG
41	ArgGluArgSerAlaAlaSerHisAlaIleThrValHisArgProLeuArgValArgGlu
	GCGGGTTTTCGCAGCCCCAGGCTTAAAGGCTGGGTAGTGGACGGTACCCCGGCCGACTGC
61	AlaGlyPheArgSerProArgLeuLysGlyTrpValValAspGlyThrProAlaAspCys
	GTCAAGCTGGGCCTGGAGGTACTTTTGCCCGAACGTCCAGATTTCCTGGTTTCGGGCATA
81	ValLysLeuGlyLeuGluValLeuLeuProGluArgProAspPheLeuValSerGlyIle
	AACTACGGGCCCAACCTGGGTACCGACGTACTTTACTCCGGCACCGTCTCGGCGGCCATA
101	AsnTyrGlyProAsnLeuGlyThrAspValLeuTyrSerGlyThrValSerAlaAlaIle
	GAAGGGGTAATTAACGGCATTCCCTCGGTGGCCGTATCTTTGGCCACGCGGCGGAGCCG
121	GluGlyValIleAsnGlyIleProSerValAlaValSerLeuAlaThrArgArgGluPro
	GACTATACCTGGGCGGCCCGGTTCGTCCTGGTCCTGCTGGAGGAACTGCGAAAACACCAA
141	AspTyrThrTrpAlaAlaArgPheValLeuValLeuLeuGluGluLeuArgLysHisGln
	CTGCCCCAGGAACCCTGCTCAACGTCAACGTGCCCGACGGGTGCCCCGCGGGGTCAAG
161	LeuProProGlyThrLeuLeuAsnValAsnValProAspGlyValProArgGlyValLys
	GTGACCAAACTGGGAAGCGTACGCTACGTCAACGTGGTAGACTGCCGCACCGACCCTCGG ValThrLysLeuGlySerValArgTyrValAsnValValAspCysArgThrAspProArg
181	
	GGGAAGGCTTACTACTGGATGGCGGGAGAACCATTGGAGCTGGACGGCAACGACTCCGAAGlyLysAlaTyrTyrTrpMetAlaGlyGluProLeuGluLeuAspGlyAsnAspSerGlu
201	
	ACCGACGTCTGGGCGGTGCGAGAAGGCTATATTTCCGTAACACCGGTCCAGATCGACCTTThrAspValTrpAlaValArgGluGlyTyrIleSerValThrProValGlnIleAspLeu
221	
	ACTAACTACGGCTTCCTGGAAGAACTCAAAAAATGGCGTTTCAAGGATATCTTTTCTTCT ThrAsnTyrGlyPheLeuGluGluLeuLysLysTrpArgPheLysAspIlePheSerSer
241	ThrashTyrGiyrheLedGidGidLedDySDySTP
261	TAA End 261
261	End 261

FIGURE 2

Methanococcus igneus Kol5 Phosphatase (9AlA) Complete Gene Sequence

	ATGTTGGATATACTGCTTGTTAATGATGATGGCATTTATTCAAATGGATTAATAGCTTTG
1	MetLeuAspIleLeuLeuValAsnAspAspGlyIleTyrSerAsnGlyLeuIleAlaLeu
	AAGGATGCATTATTGGAAAAATTTAATGCGAGGATTACTATTGTAGCCCCAACAAATCAG
21	LysAspAlaLeuLeuGluLysPheAsnAlaArgIleThrIleValAlaProThrAsnGln
	CAGAGTGGTATTGGTAGGGCAATAAGTTTATTCGAGCCGTTAAGGATAACTAAAACCAAA
41	GlnSerGlyIleGlyArgAlaIleSerLeuPheGluProLeuArgIleThrLysThrLys
	TTAGCAGATGGTTCTTGGGGATATGCAGTTTCAGGAACCCCAACAGATTGCGTTATATTG
61	LeuAlaAspGlySerTrpGlyTyrAlaValSerGlyThrProThrAspCysValIleLeu
	GGCATTTATGAGATATTAAAGAAGGTACCTGATGTAGTTATATCAGGAATAAACATTGGA
81	GlyIleTyrGluIleLeuLysLysValProAspValValIleSerGlyIleAsnIleGly
	GAAAACCTTGGGACTGAAATAACAACTTCTGGAACGTTGGGGGCTGCGTTTGAAGGGGCC
101	GluAsnLeuGlyThrGluIleThrThrSerGlyThrLeuGlyAlaAlaPheGluGlyAla
	CATCATGGGGCTAAGGCATTAGCATCACCCCAAGTTACCTCTGACCATCTAAAGTTT
121	HisHisGlyAlaLysAlaLeuAlaSerSerLeuGlnValThrSerAspHisLeuLysPhe
	AAAGAGGGGGAGACCCCAATAGACTTCACAGTCCCAGCAAGAATTACTGCAAATGTTGTT
141	LysGluGlyGluThrProIleAspPheThrValProAlaArgIleThrAlaAsnValVal
	GAGAAGATGTTGGATTATGATTTCCCATGTGATGTCGTCAACTTAAACATTCCAGAAGGA
161	GluLysMetLeuAspTyrAspPheProCysAspValValAsnLeuAsnIleProGluGly
	GCAACAGAAAAGACACCGATTGAAATCACAAGGTTGGCAAGGAAAATGTATACAACACAC
181	AlaThrGluLysThrProIleGluIleThrArgLeuAlaArgLysMetTyrThrThrHis
	GTTGAGGAAAGAATAGATCCAAGAGGGAGGAGTTATTATTGGATTGATGGGTATCCTATT
201	ValGluGluArgIleAspProArgGlyArgSerTyrTyrTrpIleAspGlyTyrProIle
	TTAGAGGAAGAGAGACACTGATGTCTATGTTGTTAGAAGAAAGGGACATATTTCTCTA
221	LeuGluGluGluHspThrAspValTyrValValArgArgLysGlyHisIleSerLeu
	ACCCCATTAACATTAGACACAACAATTAAAAATTTAGAGGAATTTAAGAAAAAATATGAG
241	ThrProLeuThrLeuAspThrThrIleLysAsnLeuGluGluPheLysLysLysTyrGlu
	AGAATATTAAATGAATGA
261	ArgIleLeuAsnGluEnd 266

FIGURE 3

Thermococcus alcaliphilus AEDII12RA Phosphatase (18A)
Complete Gene Sequence

1	ATGATGATGGAATTCACTCGCGAGGGAATAAAAGCTGCTGTAGAGGCACTTCAAGGGTT. MetMetMetGluPheThrArgGluGlyIleLysAlaAlaValGluAlaLeuGlnGlyLe
	GGAGAGATCTACGTAGTTGCCCCAATGTTTCAAAGGAGCGCAAGTGGAAGGGCAATGAC
21	GlyGluIleTyrValValAlaProMetPheGlnArgSerAlaSerGlyArgAlaMetTh
41	ATCCACAGACCTCTAAGGGCTAAAAGAATAAGTATGAACGGTGCAAAAGCAGCCTATGCTIleHisArgProLeuArgAlaLysArgIleSerMetAsnGlyAlaLysAlaAlaTyrAla
	TTGGATGGAATGCCCGTTGATTGCGTTATCTTTGCCATGGCCAGATTTGGAGATTTCGA
61	LeuAspGlyMetProValAspCysValIlePheAlaMetAlaArgPheGlyAspPheAsp
0.1	CTTGCAATAAGTGGTGTAAACTTGGGAGAAAACATGAGCACCGAGATAACGGTTTCCGGCLeuAlaIleSerGlyValAsnLeuGlyGluAsnMetSerThrGluIleThrValSerGly
81	
101	ACTGCAAGCGCTGCAATAGAGGCTGCAACCCAAGAGATCCCAAGCATTCCCATAAGCCTCThrAlaSerAlaAlaIleGluAlaAlaThrGlnGluIleProSerIleProIleSerLeu
101	IMMIASEIMIAMIAITEGIUMIAMIAIMIGIMGIAITETTOSEITIETTOSEISE
	GAAGTTAATAGAGAAAAACACAAATTTGGTGAGGGCGAAGAGATTGACTTCTCAGCTGCC
121	GluValAsnArgGluLysHisLysPheGlyGluGlyGluGluIleAspPheSerAlaAla
	AAGTATTTCCTAAGAAAAATCGCAACGGCGGTTTTAAAGAGAGGCCTCCCCAAAGGAGTC
141-	LysTyrPheLeuArgLysIleAlaThrAlaValLeuLysArgGlyLeuProLysGlyVal
	GATATGCTGAACGTCAACGTCCCTTATGATGCAAATGAAAGGACAGAGATAGCTTTTACT
161	AspMetLeuAsnValAsnValProTyrAspAlaAsnGluArgThrGluIleAlaPheThr
	CGCCTGGCAAGAAGGATGTATAGGCCTTCTATTGAAGAGCGCATAGACCCAAAGGGGAAT
181-	ArgLeuAlaArgArgMetTyrArgProSerIleGluGluArgIleAspProLysGlyAsr
	CCCTACTACTGGATAGTTGGAACTCAGTGCCCTAAGGAGGCATTAGAGCCGGGAACGGAT
201	ProTyrTrpIleValGlyThrGlnCysProLysGluAlaLeuGluProGlyThrAsp
	ATGTATGTAGTTAAAGTTGAGAGAAAAGTTAGCGTGACTCCAATAAACATTGATATGACA
221	${\tt MetTyrValValLysValGluArgLysValSerValThrProIleAsnIleAspMetThr}$
	GCAAGAGTGAATTTAGACGAGATTAAAAGACTTTTAGAACTGTAG
241	AlaArgValAsnLeuAspGluIleLysArgLeuLeuGluLeuEnd 255

FIGURE 4

Thermococcus celer Phosphatase (25A1A) Complete Gene Sequence

	ATGAGAACCCTGACAATAAACACTGACGCGGAGGGGTTCGTTTTGAGGATTCTCCTGACG	
1	MetArgThrLeuThrIleAsnThrAspAlaGluGlyPheValLeuArgIleLeuLeuThr	20
	AACGACGATGGAATCTACTCCAACGGACTGCGCGCCGCTGTGAAAGCCCTGAGTGAG	
21	AsnAspAspGlyIleTyrSerAsnGlyLeuArgAlaAlaValLysAlaLeuSerGluLeu	40
	GGCGAAGTTTACGTCGTTGCCCCCCTCTTCCAGAGGAGCGCGAGCGGCAGGGCCATGACG	
41	GlyGluValTyrValValAlaProLeuPheGlnArgSerAlaSerGlyArgAlaMetThr	60
	CTCCACAGGCCGATAAGGGCCAAGCGCGTTGACGTTCCCGGCGCAAAGATAGCCTACGGA	
61	LeuHisArgProIleArgAlaLysArgValAspValProGlyAlaLysIleAlaTyrGly	80
	ATAGATGGAACTCCTACTGACTGCGTGATTTTCGCCATAGCCCGCTTCGGGAGCTTTGGT	
81	IleAspGlyThrProThrAspCysValllePheAlaIleAlaArgPheGlySerPheGly	100
	TTAGCCGTGAGCGGGATTAACCTCGGCGAGAACCTGAGCACCGAGATAACAGTCTCAGGG	
101	LeuAlaValSerGlyIleAsnLeuGlyGluAsnLeuSerThrGluIleThrValSerGly	120
	ACGGCCTCCGCTGCCATAGAGGCCTCAACTCATGGAATTCCGAGCATAGCGATTAGCCTT	
121	ThrAlaSerAlaAlaIleGluAlaSerThrHisGlyIleProSerIleAlaIleSerLeu	140
	GAGGTGGAGTGGAAGAAGACCCTCGGCGAGGGTGAGGGGGTTGACTTCTCGGTCTCGACT	
141	GluValGluTrpLysLysThrLeuGlyGluGlyGluGlyValAspPheSerValSerThr	160
	CACTTCCTCAAGAGAATCGCGGGAGCCCTCTTGGAGAGAGGTCTTCCTGAGGGCGTTGAC	
161	HisPheLeuLysArgIleAlaGlyAlaLeuLeuGluArgGlyLeuProGluGlyValAsp	180
	ATGCTCAACGTCAACGTTCCGAGCGACGCGACGGAGGAAACGGAGATAGCAATCACCCGC	
181	MetLeuAsnValAsnValProSerAspAlaThrGluGluThrGluIleAlaIleThrArg	200
	TTAGCCCGGAAGCGCTACTCCCCAACGGTCGAGGAGGAGTTGACCCCAAGGGCAACCCC	
201	LeuAlaArgLysArgTyrSerProThrValGluGluArgIleAspProLysGlyAsnPro	220
	TACTACTGGATTGTCGGCAAACTTGTCCAAGACTTCGAGCCAGGGACAGATGCCTACGCC	
221	TyrTyrTrpIleValGlyLysLeuValGlnAspPheGluProGlyThrAspAlaTyrAla	240
	CTGAAGGTCGAGAGGAAGGTCAGGCGTCACGCCGATAAACATAGATATGACTGCGAGGGTG	
241	LeuLysValGluArgLysValSerValThrProIleAsnIleAspMetThrAlaArgVal	260
	GACTTTGAGGAGCTTGTAAGGGTTCTGTGGGTGTAA	
261	AspPheGluGluLeuValArgValLeuTrpValEnd 272	

FIGURE 5A

Thermococcus GU5L5 Phosphatase (26A1A) Complete Gene Sequence (Part 1 of 2)

	ATGAAAGGAAAGTCTCTTGTTAGCGGTCTGTTGTTGGGTCTTTTAATTTTGAGCCTGATT	
1	MetLysGlyLysSerLeuValSerGlyLeuLeuGlyLeuLeuIleLeuSerLeuIle	20
	TCATTCCAGCCAAGCTTTGCATACTCCCCACACGGCGGTGTCAAAAACATCATAATCCTG	
21	SerPheGlnProSerPheAlaTyrSerProHisGlyGlyValLysAsnIleIleIleLeu	40
	GTTGGAGACGGCATGGGTCTTGGGCATGTAGAAATTACAAAGCTCGTTTATGGACACTTA	
41	ValGlyAspGlyMetGlyLeuGlyHisValGluIleThrLysLeuValTyrGlyHisLeu	60
	AACATGGAAAACTTTCCAGTTACTGGATTTGAGCTTACTGATTCCCTAAGTGGTGAAGTT	
61	AsnMetGluAsnPheProValThrGlyPheGluLeuThrAspSerLeuSerGlyGluVal	80
	ACAGATTCTGCTGCGGCAGGAACTGCAATATCCACTGGAGCTAAAACGTATAATGGTATG	
81	ThrAspSerAlaAlaAlaGlyThrAlaIleSerThrGlyAlaLysThrTyrAsnGlyMet	100
	ATTTCAGTAACCAACATAACCGGAAAGATAGTTAACTTAACAACCCTACTTGAAGTGGCT	
101	IleSerValThrAsnIleThrGlyLysIleValAsnLeuThrThrLeuLeuGluValAla	120
	CAAGAGCTTGGGAAGTCAACAGGGCTGGTCACCACAACAAGGATTACCCATGCAACTCCA	
121	GlnGluLeuGlyLysSerThrGlyLeuValThrThrThrArgIleThrHisAlaThrPro	140
	GCAGTTTTTGCGTCCCATGTCCCAGATAGGGATATGGAGGGGGAGATACCCAAGCAACTC	
141	AlaValPheAlaSerHisValProAspArgAspMetGluGlyGluIleProLysGlnLeu	160
	ATAATGCACAAAGTTAACGTCTTGTTGGGTGGTGGAAGGGAGAAATTCGATGAGAAAAAT	
161	IleMetHisLysValAsnValLeuLeuGlyGlyGlyArgGluLysPheAspGluLysAsn	180
	TTGGAGCTGGCCAAAAAGCAGGGATACAAAGTAGTTTTCACGAAGGAAG	
181	LeuGluLeuAlaLysLysGlnGlyTyrLysValValPheThrLysGluGluLeuGluLys	200
	GTTGAAGGAGATTATGTCCTAGGACTCTTTGCAGAAAGTCACATCCCTTACGTATTGGAT	
201	ValGluGlyAspTyrValLeuGlyLeuPheAlaGluSerHisIleProTyrValLeuAsp	220
-	AGAAAACCCGATGATGTTGGACTTTTAGAAATGGCCAAAAAGGCAATTTCAATACTCGAG	246
221	ArgLysProAspAspValGlyLeuLeuGluMetAlaLysLysAlaIleSerIleLeuGlu	240
	AAGAACCCGAGCGGATTCTTTCTCATGGTTGAGGGCGGAAGGATTGACCATGCAGCCCAT	
241	LysAsnProSerGlyPhePheLeuMetValGluGlyGlyArgIleAspHisAlaAlaHis	260
	GGAAACGATGTCGCATCGGTTGTTGCAGAAACTAAGGAGTTTGACGATGTTGTCAGATAC	280
261	GlyAsnAspValAlaSerValValAlaGluThrLysGluPheAspAspValValArgTyr	200
	GTGCTGGAATATCCGAAGAAGAGGGGAGATACCTTGGTAATAGTGCTTGCCGATCACGAA	
81	ValLeuGluTyrProLysLysArgGlyAspThrLeuValIleValLeuAlaAspHisGlu	300
	ACTGGAGGTCTTGCAATAGGTCTAACGTATGGAAATGCAATCGATGAAGATGCCATAAGA	220
01	ThrGlyGlyLeuAlaIleGlyLeuThrTyrGlyAsnAlaIleAspGluAspAlaIleArg	320
	AAAATAAAAGCAAGCACCTTGAGGATGCCCAAAGAGGTTAAGGCAGGGAGTAGTGTAAAA	
21	LystleLysAlaSerThyLeuArgMetProLysGluValLysAlaGlySerSerValLys	340

FIGURE 5B

Thermococcus GU5L5 Phosphatase (26A1A) Complete Gene Sequence (Part 2 of 2)

	GAGTCCTCAAAGGTATGCCGGATTTGTCCCAACAGAGGAAGTCAGTATATTGAGAAT	
341	GluSerSerLysValCysArgIleCysProAsnArgGlyArgSerGlnTyrIleGluAsn	360
	GCGCTGCACTCGACAAACAAGTATGCCCTCTCAAATGCAGTAGCCGATGTTATAAACAGG	
361	AlaLeuHisSerThrAsnLysTyrAlaLeuSerAsnAlaValAlaAspValIleAsnArg	380
	CGTATTGGTGTTGGATTCACCTCCTATGAGCATACAGGAGTTCCAGTTCCGCTCTTAGCT	
381	ArgIleGlyValGlyPheThrSerTyrGluHisThrGlyValProValProLeuLeuAla	400
	TACGGTCCCGGGCAGAGAACTTCAGAGGTTTCTTACACCATGTGGATACAGCAAGATTA	
401	TyrGlyProGlyAlaGluAsnPheArgGlyPheLeuHisHisValAspThrAlaArgLeu	420
	GTTGCAAAGTTAATGCTCTTTGGAAGGAGGAATATTCCAGTTACCATTTCAAGCGTGAGC	
421	ValAlaLysLeuMetLeuPheGlyArgArgAsnIleProValThrIleSerSerValSer	440
	AGTGTTAAGGGAGACATAACCGGTGATTACAGGGTTGATGAGAAGGATGCCTACGTTACG	
441	SerValLysGlyAspIleThrGlyAspTyrArgValAspGluLysAspAlaTyrValThr	460
	CTCATGATGTTTCTCGGAGAAAAAGTGGATAATGAAAATTGAAAAGAGAGTCGATATAGAC	
461	LeuMetMetPheLeuGlyGluLysValAspAsnGluIleGluLysArgValAspIleAsp	480
	AACAACGGCATGGTTGACTTAAATGACGTCATGTTGATTCTCCAGGAAGCTTGA	
481	AsnAsnGlyMetValAspLeuAsnAspValMetLeuIleLeuGlnGluAlaEnd 498	

FIGURE 6A

OC9a Phosphatase (27A3A) Complete Gene Sequence (Part 1 of 2)

	ATGCCAAGAAATATCGCCGCTGTATGCGCCCTTGGCCGCTTTGTTAGGGTCGGCCTGGGCG	
1	MetProArgAsnIleAlaAlaValCysAlaLeuAlaAlaLeuLeuGlySerAlaTrpAla	20
	GCCAAAGTTGCCGTCTACCCCTACGACGGAGCCGCTTTGCTGGCGGGGGGGCAGCGCTTCGAT	
21	AlaLysValAlaValTyrProTyrAspGlyAlaAlaLeuLeuAlaGlyGlnArgPheAsp	40
	TTGCGCATAGAAGCCTCCGAGCTGAAAGGCAATTTAAAGGCTTACCGCATCACCCTGGAC	
41	LeuArgIleGluAlaSerGluLeuLysGlyAsnLeuLysAlaTyrArgIleThrLeuAsp	60
	GGCCAGCCTCTGGCGGCCTCGAGCAAACCGCGCAGGGGGCCGGGCAGGCCGAGTGGACC	
61	${\tt GlyGlnProLeuAlaGlyLeuGluGlnThrAlaGlnGlyAlaGlyGlnAlaGluTrpThr}$	80
	CTGCGCGTGCCTTCCTGCGCCCTGGAAGCCACACCCTCGAGGTCAGCCTCACCGACGAC	
81	LeuArgGlyAlaPheLeuArgProGlySerHisThrLeuGluValSerLeuThrAspAsp	100
91	Leuki golyklarnebeuki gri oolybeliila imaabaata valbel bealiilkaspaap	100
	GCTGGGGAGAGCAGGAAGAGCGTACGTTGGGAGGCTCGGCAGAACCTTCGCTTGCCCCGA	
101	AlaGlyGluSerArgLysSerValArgTrpGluAlaArgGlnAsnLeuArgLeuProArg	120
	GCGGCCAAGAATGTGATTCTCTTCATTGGCGACGGGATGGGCTGGAACACCCTCAACGCC	
121	${\tt AlaAlaLysAsnValIleLeuPheIleGlyAspGlyMetGlyTrpAsnThrLeuAsnAla}$	140
	GCCCGCATCATCGCCAAAGGCTTTAACCCCGAAAACGGTATGCCCAACGGAAACCTCGAG	
	AlaArgIleIleAlaLysGlyPheAsnProGluAsnGlyMetProAsnGlyAsnLeuGlu	160
141	AlaargiteiteataLysgiyPheashFlootuashotyMetrioAshGiyAshBedGid	100
	ATCGAGAGTGGTTACGGTGGGATGGCTACCGTCACTACCGGCAGCTTTGATAGCTTCATC	
161	${\tt IleGluSerGlyTyrGlyGlyMetAlaThrValThrThrGlySerPheAspSerPheIle}$	180
	GCCGACTCAGCTAACTCGGCTTCTTCCATCATGACCGGGCAGAAGGTGCAGGTGAATGCC	
181	$\verb AlaAspSerAlaAsnSerAlaSerSerIleMetThrGlyGlnLysValGlnValAsnAla $	200
	CTCAACGTTTACCCATCAAACCTCAAAGATACCCTGGCCTACCCCGGATCGAAACCCTA	
201	LeuAsnValTyrProSerAsnLeuLysAspThrLeuAlaTyrProArgIleGluThrLeu	220
201	Demy20.411.11.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.11.12.1	
	GCGGAGATGCTCAAGCGGGTACGCGGGGCCAGCATTGGGGTAGTGACCACCACCTTCGGC	
221	AlaGluMetLeuLysArgValArgGlyAlaSerIleGlyValValThrThrThrPheGly	240
	ACCGACGCTACCCCGGCTTCACTCAACGCCCATACCCGCCGCCGCGGTGATTACCAGGCT	
241	$Thr {\tt AspAlaThrProAlaSerLeuAsnAlaHisThrArgArgArgGlyAspTyrGlnAla}$	260
	ATCGCCGACATGTACTTTGGTAGAGGCGGGTTCGGTGTTCCCTTGGATGTGATGCTCTTC	
261	IleAlaAspMetTyrPheGlyArgGlyGlyPheGlyValProLeuAspValMetLeuPhe	280
	GGTGGTTCACGCGACTTCATCCCCCAGAGCACCCCTGGCTCGCGCGCAAGGATAGCACG	200
281	GlyGlySerArgAspPheIleProGlnSerThrProGlySerArgArgLysAspSerThr	300
	GACTGGATTGCCGAATCCCAGAAGCTGGGCTACACCTTTGTCAGCACCCGCAGCGAGCTG	
01	AspTrpIleAlaGluSerGlnLysLeuGlyTyrThrPheValSerThrArgSerGluLeu	320
	CTGGCGGCCAAACCCACCGATAAGCTGTTTGGGCTGTTCAACATTGACAACTTCCCCAGC	
121	LeuAlaAlaLysProThrAspLysLouPheGlyLeuPheAsnIleAspAsnPheProSer	340
1 Z I	DEGUTAVIADADI I O LIII USBULLA INGGA II GA III GA II GA III GA II GA I	

FIGURE 6B

OC9a Phosphatase (27A3A) Complete Gene Sequence (Part 2 of 2)

341	TACCTAGACCGCGCAGTGTGGAAGCGGCCCGAGATGCTGGGAAGCTTTACCGATATGCCC TyrLeuAspArgAlaValTrpLysArgProGluMetLeuGlySerPheThrAspMetPro	360
361	thm:thm:thm:thm:thm:thm:thm:thm:thm:thm:	380
381	TTCTTGATGGTTGAGGGGGGAATGGTGGATAAGTACGAGCACCCCTTGGACTGGCCCCGC PheLeuMetValGluGlyGlyMetValAspLysTyrGluHisProLeuAspTrpProArg	400
401	GCACTTTGGGATGTACTCGAGCTGGACCGCGCGCGGTGGCTTGGGCCAAGGGCTATGCGGCC AlaLeuTrpAspValLeuGluLeuAspArgAlaValAlaTrpAlaLysGlyTyrAlaAla	420
421	TCCCACCCCGATACCCTGGTGATTGTCACCGCCGACCACGCTCACTCGATCTCGGTGTTT SerHisProAspThrLeuVallleValThrAlaAspHisAlaHisSerIleSerValPhe	440
441	GGCGGTTACGACTACTCCAAGCAGGCCGGGAGGGGGTGGGGGTTTATGAGGCCGCCAAG GlyGlyTyrAspTyrSerLysGlnGlyArgGluGlyValGlyValTyrGluAlaAlaLys	460
461	TTCCCCACCTACGGCGACAAAAAAGACGCCAACGGCTTTCCCTTGCCCGACACCACTCGG PheProThrTyrGlyAspLysLysAspAlaAsnGlyPheProLeuProAspThrThrArg	480
481	GGAATCGCGGTAGGCTTCGGGGCCACGCCGGATTACTGTGAAACCTACCGGGGCCGCGAG GlyIleAlaValGlyPheGlyAlaThrProAspTyrCysGluThrTyrArgGlyArgGlu	500
501	GTCTACAAAGACCCCACCATCTCCGACGGCAAAGGTGGTTACGTGGCCAACCCTGAGGTC ValTyrLysAspProThrIleSerAspGlyLysGlyGlyTyrValAlaAsnProGluVal	520
521	TGCAAGGAGCCGGGCCTTCCAACGTACCGGCAACTCCCAGTAGATAGCGCCCAGGGCGTG CysLysGluProGlyLeuProThrTyrArgGlnLeuProValAspSerAlaGlnGlyVal	540
541	CACACGGCTGATCCCATGCCGCTGTTTGCCTTTGGCGTGGGGTCTCAGTTCTTCAATGGC HisThrAlaAspProMetProLeuPheAlaPheGlyValGlySerGlnPhePheAsnGly	560
561	CTCATCGACCAGACCGAGATCTTCTTCCGCATGGCCCAGGCCCTAGGGTTCAACCCCCAC LeuIleAspGlnThrGluIlePhePheArgMetAlaGlnAlaLeuGlyPheAsnProHis	580
581	CTCGAGAAGCCTTAA	

FIGURE 7

Mll TL Phosphatase (29AlA=29A2A) Complete Gene Sequence

	ATGTATAAATGGATTATTGAGGGTAAGCTTGCCCAAGCACCTTTTCCAAGCCTAGGTGAA	2.0
1	MetTyrLysTrpIleIleGluGlyLysLeuAlaGlnAlaProPheProSerLeuGlyGlu	20
	CTAGCCGATCTCAAAAGACTTTTCGACGCCATTATTGTTCTTACAATGCCGCATGAACAA	
21	LeuAlaAspLeuLysArgLeuPheAspAlaIleIleValLeuThrMetProHisGluGln	40
	CCGCTTAATGAGAAATATATCGAGATATTAGAGAGCCATGGATTCCAAGTCCTCCATGTC	
41	ProLeuAsnGluLysTyrIleGluIleLeuGluSerHisGlyPheGlnValLeuHisVal	60
	CCCACGCTCGACTTTCATCCTTTAGAACTCTTCGACCTTTTGAAAACAAGCATATTCATT	
61	ProThrLeuAspPheHisProLeuGluLeuPheAspLeuLeuLysThrSerIlePheIle	80
	GATGAAAACCTGGAGAGATCCCACAGAGTGCTTGTCCACTGCATGGGAGGCATAGGCCGG	
81	AspGluAsnLeuGluArgSerHisArgValLeuValHisCysMetGlyGlyIleGlyArg	100
	AGCGGGCTTGTAACTGCTGCGTACTTAATATTCAAAGGTTATGATATTTACGACGCGGTA	
101	SerGlyLeuValThrAlaAlaTyrLeuIlePheLysGlyTyrAspIleTyrAspAlaVal	120
	AAGCATGTGAGAACGGTAGTGCCTGGTGCTATTGAAAACAGAGGGCAAGCGTTAATGCTT	
121	LysHisValArgThrValValProGlyAlaIleGluAsnArgGlyGlnAlaLeuMetLeu	140
	GAGAACTACTATACCCTGGTCAAAAGTTTCAACAGAGAGTTGCTGAGAGACTACGGGAAG	
141	GluAsnTyrTyrThrLeuValLysSerPheAsnArgGluLeuLeuArgAspTyrGlyLys	160
	AAAATTTTCACGCTCGGTGACCCGAAGGCGGTTCTCCACGCTTCTAAGACGACTCAGTTC	
161	LysIlePheThrLeuGlyAspProLysAlaValLeuHisAlaSerLysThrThrGlnPhe	180
	ACGATTGAACTCTTAAGCAACTTACACGTCAACGAGGCGTTTTCAATCAGTGCGATGGCT	
181	ThrIleGluLeuLeuSerAsnLeuHisValAsnGluAlaPheSerIleSerAlaMetAla	200
	CAATCACTGCTCCACTTTCACGACGTAAAAGTCCGCTCTAAACTGAAAGAAGTATTCGAA	
201	GlnSerLeuLeuHisPheHisAspValLysValArgSerLysLeuLysGluValPheGlu	220
	AACATGGAATTCTCATCCGCCTCAGAGGAGGTTCTGTCATTTATTCACCTACTCGATTTC	240
221	AsnMetGluPheSerSerAlaSerGluGluValLeuSerPheIleHisLeuLeuAspPhe	240
	TATCAGGATGGCAGGGTTGTTTTAACCATTTACGATTATCTCCCCGATAGGGTGGATTTG	260
241	TyrGlnAspGlyArgValValLeuThrIleTyrAspTyrLeuProAspArgValAspLeu	260
	ATTTTATTGTGTAAGTGGGGTTGTGATAAAATAGTTGAAGTCTCGTCTTCAGCGAAGAAA	200
261	IleLeuLeuCysLysTrpGlyCysAspLysIleValGluValSerSerSerAlaLysLys	280
	ACCGTTGAGAAGCTTGTAGGAAGAAGGTTTCCCTATCCTGGGCTAATTACTTAGACTAT	300
281	ThrValGluLysLeuValGlyArgLysValSerLeuSerTrpAlaAsnTyrLeuAspTyr	ייינ
	GTTTTAG	
301	ValEnd 102	

FIGURE 8

Thermococcus CL-2 Phosphatase (30A1A) Complete Gene Sequence

	ATGAGAATCCTCCTCACCAACGACGACGGCATCTATTCCAACGGTCTGCGCGCGGCGGTG	
1	MetArgIleLeuLeuThrAsnAspAspGlyIleTyrSerAsnGlyLeuArgAlaAlaVal	20
	AAGGGCCTGAGCGAGCTCGGCGAGGTCTACGTCGTCGCCCCGCTCTTCCAGAGGAGCGCG	
21	LysGlyLeuSerGluLeuGlyGluValTyrValValAlaProLeuPheGlnArgSerAla	40
	AGCGGTCGGGCGATGACCCTACACAGGCCGATAAGGGCAAAGAGGGTTGACGTTCCCGGC	
41	SerGlyArgAlaMetThrLeuHisArgProIleArgAlaLysArgValAspValProGly	60
	GCGAAGATAGCGTATGGCATAGACGGAACGCCGACCGACTGCGTGATTTTTGCCATCGCC	•
61	AlaLysIleAlaTyrGlyIleAspGlyThrProThrAspCysValIlePheAlaIleAla	80
	CGCTTCGGCGACTTTGATCTGGCGGTCAGCGGGATAAACCTAGGCGAGAACCTGAGCACG	
81	ArgPheGlyAspPheAspLeuAlaValSerGlyIleAsnLeuGlyGluAsnLeuSerThr	100
	GAGATAACCGTCTCCGGAACGGCCTCGGCGGCGATAGAGGCTTCCACCCAC	
101	GlulleThrValSerGlyThrAlaSerAlaAlaIleGluAlaSerThrHisGlyIlePro	120
	AGTGTAGCTATAAGCCTCGAGGTCGAGTGGAAGAAGACCCTCGGCGAGGGGAGGGTATT	
121	SerValAlaIleSerLeuGluValGluTrpLysLysThrLeuGlyGluGlyGluGlyIle	140
	GACTTCTCGGTTTCAGCACACTTCCTGAGAAGGATAGCGACGGCTGTCCTTAAGAAGGGC	
141	AspPheSerValSerAlaHisPheLeuArgArgIleAlaThrAlaValLeuLysLysGly	160
	CTGCCTGAAGGGGTGGACATGCTCAACGTGAACGTCCCTAGCGACGCCAGCGAGGGGACT	
161	LeuProGluGlyValAspMetLeuAsnValAsnValProSerAspAlaSerGluGlyThr	180
	GAGATCGCCATAACGCGCCTCGCGAGGAAGCGCTATTCTCCGACGATAGAGGAGAGGATA	
181	GluIleAlaIleThrArgLeuAlaArgLysArgTyrSerProThrIleGluGluArgIle	200
	GACCCCAAGGGCAACCCCTACTACTGGATCGTTGGCAGGCTCGTCCAGGAGTTCGAGCCG	
201	AspProLysGlyAsnProTyrTyrTrpIleValGlyArgLeuValGlnGluPheGluPro	220
	GGCACGGACGCCTACGCTCTGAAAGTCGAGAGAAAGGTCAGCGTCACGCCCATAAACATC	
221	GCACGGACGCCTACGCTCTGABACTCTGACGCTGACACGCTGACGACGACGCTGACACGCTGACACACCACACACA	240
	GACATGACTGCGAGGGTTGACTTTGAGAACCTTCAAAGGCTTCTGAGCCTGTGA	
243	AspMotThralaArgValAspPheGluAsnLeuGlnArgLeuLeuSerLeuEnd 258	

11/11

FIGURE 9

Aquifex VF-5 Phosphatase (34A1A) Complete Gene Sequence

	ATGGAAAACTTAAAAAAGTACCTAGAAGTTGCAAAAATAGCCGCGCTCGCGGGTGGGCAG	2.0
1	MetGluAsnLeuLysLysTyrLeuGluValAlaLysIleAlaAlaLeuAlaGlyGlyGln	20
	GTTCTGAAAGAAACTTCGGAAAGGTAAAAAAGGAAAACATAGAGGAAAAAGGGGAAAAAG	
21	ValLeuLysGluAsnPheGlyLysValLysLysGluAsnIleGluGluLysGlyGluLys	40
	GACTTTGTAAGTTACGTGGATAAAACTTCAGAGGAAAGGATAAAGGAGGTGATACTCAAG	
41	AspPheValSerTyrValAspLysThrSerGluGluArgIleLysGluValIleLeuLys	60
	TTCTTTCCCGATCACGAGGTCGTAGGGGAAGAGATGGGTGCGGAGGGAAGCGGAAGCGAA	
61	PhePheProAspHisGluValValGlyGluGluMetGlyAlaGluGlySerGlySerGlu	80
	TACAGGTGGTTCATAGACCCCCTTGACGGCACAAAGAACTACATAAACGGTTTTCCCATC	
81	TyrArgTrpPheIleAspProLeuAspGlyThrLysAsnTyrIleAsnGlyPheProIle	100
	TTTGCCGTATCAGTGGGACTTGTTAAGGGAGAAGAGCCAATTGTGGGTGCGGTTTACCTT	
101	PheAlaValSerValGlyLeuValLysGlyGluGluProIleValGlyAlaValTyrLeu	120
	CCTTACTTTGACAAGCTTTACTGGGGTGCTAAAGGTCTCGGGGCTTACGTAAACGGAAAG	
121	ProTyrPheAspLysLeuTyrTrpGlyAlaLysGlyLeuGlyAlaTyrValAsnGlyLys	140
	AGGATAAAGGTAAAGGACAATGAGAGTTTAAAGCACGCCGGAGTGGTTTACGGATTTCCC	
141	ArgIleLysValLysAspAsnGluSerLeuLysHisAlaGlyValValTyrGlyPhePro	160
	TCTAGGAGCAGGAGGACATATCTATCTACTTGAACATATTCAAGGATGTCTTTTACGAA	
161	SerArgSerArgArgAspIleSerIleTyrLeuAsnIlePheLysAspValPheTyrGlu	180
	GTTGGCTCTATGAGGAGACCCGGGGCTGCTGCGGTTGACCTCTGCATGGTGGCGGAAGGG	
181	ValGlySerMetArgArgProGlyAlaAlaAlaValAspLeuCysMetValAlaGluGly	200
	ATATTTGACGGGATGATGGAGTTTGAAATGAAGCCGTGGGACATAACCGCAGGGCTTGTA	
201	IlePheAspGlyMetMetGluPheGluMetLysProTrpAspIleThrAlaGlyLeuVal	220
	ATACTGAAGGAAGCCGGGGGCGTTTACACACTTGTGGGAGAACCCTTCGGAGTTTCGGAC	
221	IleLeuLysGluAlaGlyGlyValTyrThrLeuValGlyGluProPheGlyValSerAsp	240
	ATAATTGCGGGCAACAAAGCCCTCCACGACTTTATACTTCAGGTAGCCAAAAAGTATATG	
241	IleIleAlaGlyAsnLysAlaLeuHisAspPheIleLeuGlnValAlaLysLysTyrMet	260
	GAAGTGGCGGTGTGA	
261	GluValAlaValEnd 265	

INTERNATIONAL SEARCH REPORT

International application No. PCT/US97/10784

		 		
A. CLASSIFICATION OF SUBJECT MATTER				
IPC(6) US CL	:A61K 38/46; C07H 19/00, 21/02, 21/04; C12N 9/14 :424/94.6; 435/95, 252.3, 320.1; 536/22.1, 23.1, 23.2			
According t	to International Patent Classification (IPC) or to both	national classification and IPC		
	DS SEARCHED			
Minimum d	ocumentation searched (classification system followe	d by classification symbols)		
U.S. :	424/94.6; 435/95, 252.3, 320.1; 536/22.1, 23.1, 23.2;	935/22		
Documental	tion searched other than minimum documentation to the	e extent that such documents are included	in the fields searched	
Electronic d	lata base consulted during the international search (na	ame of data base and, where practicable,	, search terms used)	
Please See	e Extra Sheet.			
C. DOC	UMENTS CONSIDERED TO BE RELEVANT			
Category*	Citation of document, with indication, where ap	propriate, of the relevant passages	Relevant to claim No.	
A	MOSS. Perspectives in Alkaline Pho Chemistry. December 1992. Vol. 38	- ,	1-12	
A	MOSS et al. Clinical and Biological Aspects of Acid Phosphatase. Critical Reviews in Clinical Laboratory Sciences. July 1995. Vol. 32, No. 4, pages 431-467.			
	_			
Furth	er documents are listed in the continuation of Box C	See patent family annex.		
	cial categories of cited documents:	"I" later document published after the unit date and not in conflict with the appl		
	nument defining the general state of the art which is not considered be of particular relevance	the principle or theory underlying the		
'B' oar	lier document published on or after the international filing date	"X" document of particular relevance; the considered novel or cannot be considered.		
cite	nument which may throw doubts on priority claim(s) or which is do to establish the publication date of another citation or other	when the document is taken alone		
special reason (as specified) "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is				
"O" document referring to an oral disclosure, use, exhibition or other combined with one or more other such documents, such combination being obvious to a person skilled in the art				
P document published prior to the international filing data but later than *a.* document member of the same patent family the priority data claimed				
Date of the actual completion of the international search Date of mailing of the international search report				
10 SEPTEMBER 1997 28 OCT 1997				
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Authorized officer				
Box PCT				
	acsimile No. (703) 305-3230 Telephone No. (703) 308-0196			

INTERNATIONAL SEARCH REPORT

International application No. PCT/US97/10784

B. FIELDS SEARCHED Electronic data bases consulted (Name of data base and where practicable terms used):
database: APS, CAS ONLINE, CAS REGISTRY, MEDLINE, BIOSIS, EMBASE, GENBANK search terms: SEQ ID NO: 19-54, ammonifex, aquifex methanococcus, thermococcus, alkaline phosphatase, thermostable
-